CITATION REPORT List of articles citing

Clinical implications of the wo reflowwphenomenon. A predictor of complications and left ventricular remodeling in reperfused anterior wall myocardial infarction.

DOI: 10.1161/01.CIR.93.2.223 Circulation, 1996, 93, 223-8.

Source: https://exaly.com/paper-pdf/86224791/citation-report.pdf

Version: 2024-04-17

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
777	New insights into the pathogenesis, prevention and management of acute coronary syndromes. 1996 , 26, 344-8		3
776	Myocardial contrast echocardiography to assess spontaneous reperfusion during myocardial infarction. 1997 , 349, 617-9		3
775	Clinical value of 12-lead electrocardiogram after successful reperfusion therapy for acute myocardial infarction. Zwolle Myocardial infarction Study Group. 1997 , 350, 615-9		459
774	Beneficial effect of intracoronary verapamil on microvascular and myocardial salvage in patients with acute myocardial infarction. 1997 , 30, 1193-9		223
773	Recovery of myocardial perfusion in acute myocardial infarction after successful balloon angioplasty and stent placement in the infarct-related coronary artery. 1997 , 30, 1270-6		66
77²	Usefulness of magnetic resonance imaging early after acute myocardial infarction. 1997 , 80, 690-5		34
771	Frequency of pericardial friction rub ("pericarditis") after direct percutaneous transluminal coronary angioplasty in Q-wave acute myocardial infarction. 1997 , 79, 362-4		2
77°	Current role of contrast echocardiography in the diagnosis of cardiovascular diseases. 1997 , 20, I31-8		6
769	Primary PTCA: Possibly the Best, Often the Only Choice for Reperfusion in Acute Myocardial Infarction. 1997 , 4, 331-336		1
768	Myocardial Reperfusion Injury: From Bench to Bedside. 1997 , 4, 35-37		
767	Regional dysfunction and ventricular remodeling in the infarcted patient. 1997 , 92, 72-4		
766	Rapid reversal of no-reflow using Abciximab after coronary device intervention. 1997 , 42, 187-90		46
765	The Open Artery Hypothesis: Past, Present, and Future. 1998 , 5, 101-112		1
764	Myocardial Perfusion Abnormalities by Intravenous Administration of the Contrast Agent NC100100 in an Experimental Model of Coronary Artery Thrombosis and Reperfusion. 1998 , 15, 731-74	10	5
763	Assessing the relation between coronary reflow and myocardial reflow. 1998 , 81, 8G-12G		33
762	Contrast echocardiography: review and future directions. 1998 , 81, 41G-48G		43
761	Role of nitric oxide in coronary arterial vasomotion and the influence of coronary atherosclerosis and its risks. 1998 , 82, 1034-9		23

(1999-1998)

760	The clinical implications of no reflow demonstrated with intravenous perfluorocarbon containing microbubbles following restoration of Thrombolysis In Myocardial Infarction (TIMI) 3 flow in patients with acute myocardial infarction. 1998 , 82, 1173-7	101
759	Increased intensity of contrast material immediately after late angioplasty of infarct-related coronary artery is associated with reduced ventricular volumes at six months. 1998 , 82, 1451-6	4
758	10 Transoesophageal echocardiography: emerging and innovative technologies. 1998, 12, 681-691	
757	Primary coronary angioplasty in acute myocardial infarction: clinical correlates of the 'no reflow' phenomenon. 1998 , 65 Suppl 1, S23-8	16
756	Critical analysis of coronary artery bypass graft surgery: a 30-year journey. 1998 , 31, 1B-63B	82
755	The pattern of alteration in flow velocity in the recanalized artery is related to left ventricular recovery in patients with acute infarction and successful direct balloon angioplasty. 1998 , 32, 338-44	27
754	Prediction of short- and intermediate-term prognoses of patients with acute myocardial infarction using myocardial contrast echocardiography one day after recanalization. 1998 , 32, 890-7	146
753	Evaluation of dynamic changes in microvascular flow during ischemia-reperfusion by myocardial contrast echocardiography. 1998 , 32, 1096-101	37
75 ²	Quantification and time course of microvascular obstruction by contrast-enhanced echocardiography and magnetic resonance imaging following acute myocardial infarction and reperfusion. 1998 , 32, 1756-64	256
75 ¹	Assessment of "microvascular no-reflow phenomenon" using technetium-99m macroaggregated albumin scintigraphy in patients with acute myocardial infarction. 1998 , 32, 898-903	110
750	Early changes in myocardial perfusion patterns after myocardial infarction: relation with contractile reserve and functional recovery. 1998 , 32, 2011-7	50
749	Integrated approach to ischemic heart disease. The one-stop shop. 1998 , 16, 267-76	27
748	Added value of contrast echocardiography in assessing myocardial viability. 1999 , 82 Suppl 3, III22-6	2
747	Acute assessment of microvascular perfusion patterns by myocardial contrast echocardiography during myocardial infarction: relation to timing and extent of functional recovery. 1999 , 81, 12-6	31
746	Combined assessment of reflow and collateral blood flow by myocardial contrast echocardiography after acute reperfused myocardial infarction. 1999 , 82, 62-7	9
745	Reperfusion revisited: beyond TIMI 3 flow. 1999 , 22, IV20-9	20
744	[Prognostic significance of ST segment change in acute myocardial infarct]. 1999 , 24, 378-88	12
743	Coronary microembolizationits role in acute coronary syndromes and interventions. 1999 , 24, 558-75	41

742	[Patency, perfusion and prognosis in acute myocardial infarct]. 1999 , 24, 421-9	8
741	Current state of thrombolytic therapy. 1999 , 1, 212-20	2
740	Early temporal changes in coronary flow velocity patterns in patients with acute myocardial infarction demonstrating the "no-reflow" phenomenon. 1999 , 84, 415-9	46
739	Myocardial "low reflow" assessed by Dy-DTPA-BMA-enhanced first-pass MR imaging in a dog model. 1999 , 9, 679-84	16
738	Predictors of stent thrombosis after primary stenting for acute myocardial infarction. 1999 , 47, 415-22	12
737	Intravenous nicorandil can preserve microvascular integrity and myocardial viability in patients with reperfused anterior wall myocardial infarction. 1999 , 33, 654-60	313
736	The significance of persistent ST elevation versus early resolution of ST segment elevation after primary PTCA. 1999 , 34, 1932-8	161
735	Primary stenting in acute myocardial infarction: influence of diabetes mellitus in angiographic results and clinical outcome. 1999 , 138, 446-55	47
734	Reperfusion injury: experimental evidence and clinical implications. 1999 , 138, S69-75	245
733	Microvascular integrity after reperfusion therapy. 1999 , 138, S76-8	27
732	Early predictors of left ventricular remodeling after acute myocardial infarction. 1999 , 138, S79-83	34
731	Recanalization versus reperfusion for myocardial survival and preservation of ventricular geometry. 1999 , 138, S89-95	2
730	Clinical applications of transpulmonary contrast echocardiography. 1999 , 137, 144-53	56
729	Differential diagnosis of left ventricular mural thrombi by myocardial contrast echocardiography: a preliminary study. 1999 , 63, 50-2	5
728	Relation between coronary thrombus and angiographic no-flow during primary angioplasty in patients with acute myocardial infarction. 1999 , 63, 849-53	17
727	The use of myocardial contrast echocardiography in clinical evaluation after myocardial infarction. 2000 , 11, 235-42	1
726	Intracoronary adenosine administered during percutaneous intervention in acute myocardial infarction and reduction in the incidence of "no reflow" phenomenon. 2000 , 51, 27-31; discussion 32	82
725	Clinical experience with SonoVue in myocardial perfusion imaging. 2000 , 17, S17-23	7

724	Contrast echocardiography: clinical utility. 2000 , 17, S5-9	5
723	Calcium signaling and oxidant stress in the vasculature. 2000 , 28, 1362-9	120
722	Achieving tissue-level perfusion in the setting of acute myocardial infarction. 2000 , 85, 39C-46C	21
721	Antiplatelet therapy in interventional cardiology: II. Glycoprotein IIb/IIIa inhibitors. 2000 , 9, 163-74	2
720	Treatment strategies for microvascular dysfunction following acute myocardial infarction. 2000 , 2, 405-10	1
719	Reversal of slow flow phenomenon during primary stenting by bail-out administration of abciximab. 2000 , 3, 35-39	5
718	Absence of preinfarction angina is associated with a risk of no-reflow phenomenon after primary coronary angioplasty for a first anterior wall acute myocardial infarction. 2000 , 75, 253-60	19
717	Myocardial infarction and left ventricular remodeling: results of the CEDIM trial. Carnitine Ecocardiografia Digitalizzata Infarto Miocardico. 2000 , 139, S124-30	20
716	Importance of microembolization and inflammation in atherosclerotic heart disease. 2000, 140, S90-102	23
715	[Coronary angioplasty in the acute myocardial infarction: in which patients is it less likely to obtain an adequate coronary reperfusion?]. 2000 , 53, 1169-76	11
714	Reperfusion syndrome: relationship of coronary blood flow reserve to left ventricular function and infarct size. 2000 , 35, 1162-9	50
713	Angiographic no-reflow phenomenon as a predictor of adverse long-term outcome in patients treated with percutaneous transluminal coronary angioplasty for first acute myocardial infarction. 2000 , 36, 1202-9	438
712	Potential of microvascular reperfusion with adjunctive pharmacological intervention: its impact on myocardial perfusion and functional outcomes in patients with acute myocardial infarction. 2001 , 61, 437-41	3
711	Course and prognostic implications of QT interval and QT interval variability after primary coronary angioplasty in acute myocardial infarction. 2001 , 37, 44-50	58
710	Shifting the open-artery hypothesis downstream: the quest for optimal reperfusion. 2001 , 37, 9-18	201
709	Postinfarctional remodeling: increased dye intensity in the myocardial risk area after angioplasty of infarct-related coronary artery is associated with reduction of ventricular volumes. 2001 , 37, 1239-45	12
708	Predictive factors for development of the no-reflow phenomenon in patients with reperfused anterior wall acute myocardial infarction. 2001 , 38, 472-7	152
707	Full-motion pulse inversion power Doppler contrast echocardiography differentiates stunning from necrosis and predicts recovery of left ventricular function after acute myocardial infarction. 2001 , 38, 1390-4	74

706	Pressure-derived collateral flow index as a parameter of microvascular dysfunction in acute myocardial infarction. 2001 , 38, 1383-9	42
705	Clinical applications of contrast echocardiography. 2001 , 141, S36-44	6
704	Corrected TIMI frame counts correlate with stenosis severity and infarct zone wall motion after thrombolytic therapy. 2001 , 141, 586-91	9
703	Direct infarct artery stenting without predilation and no-reflow in patients with acute myocardial infarction. 2001 , 142, 684-90	66
702	ST segment resolution as a tool for assessing the efficacy of reperfusion therapy. 2001 , 38, 1283-94	212
701	A randomized double-blind trial of intravenous trimetazidine as adjunctive therapy to primary angioplasty for acute myocardial infarction. 2001 , 77, 263-73	41
700	Intracoronary electrocardiogram for early detection of myocardial viability during coronary angioplasty in acute myocardial infarction. 2001 , 79, 293-9	10
699	Extent of ST-segment deviation in a single electrocardiogram lead 90 min after thrombolysis as a predictor of medium-term mortality in acute myocardial infarction. 2001 , 358, 1479-86	99
698	Assessment of no-reflow phenomenon after acute myocardial infarction with harmonic angiography and intravenous pump infusion with Levovist: comparison with intracoronary contrast injection. 2001 , 14, 773-81	22
697	Combined revascularization strategy for acute myocardial infarction in patients with intracoronary thrombus: preceding intracoronary thrombolysis and subsequent mechanical angioplasty. 2001 , 65, 251-6	18
696	Prediction of wall motion recovery from the left anterior descending coronary artery velocity pattern recorded by transthoracic doppler echocardiography in patients with anterior wall myocardial infarction retrospective and prospective studies. 2001 , 65, 717-22	11
695	Pulse infusion thrombolysis (PIT) for large intracoronary thrombus: preventive effect against the 'no flow' phenomenon in revascularization therapy for acute myocardial infarction. 2001 , 65, 94-8	14
694	Evolution in the practice of primary angioplasty: effect of adjunctive coronary stenting and glycoprotein IIb/IIIa inhibitors on long-term outcomes. 2001 , 54, 327-32	6
693	Predictors and prognosis of suboptimal coronary blood flow after primary coronary angioplasty in patients with acute myocardial infarction. 2001 , 88, 124-8	69
692	Association of noninvasive markers of coronary artery reperfusion to assess microvascular obstruction in patients with acute myocardial infarction treated with primary angioplasty. 2001 , 88, 342-6	23
691	Early identification of impaired myocardial reperfusion with serial assessment of ST segments after percutaneous transluminal coronary angioplasty during acute myocardial infarction. 2001 , 88, 956-9	28
690	Predictive value of markers of myocardial reperfusion in acute myocardial infarction for follow-up left ventricular function. 2001 , 88, 1358-63	40
689	Interventions in acute myocardial infarction. 2001 , 26, 619-72	3

688 Cellular Mechanisms of Cardioprotection. 2001, 853-866

687	Angiographic morphologic features of infarct-related arteries and timely reperfusion in acute	
	myocardial infarction: predictors of slow-flow and no-reflow phenomenon. 2002 , 122, 1322-32	227
686	Can coronary flow velocity reserve determined by transthoracic Doppler echocardiography predict the recovery of regional left ventricular function in patients with acute myocardial infarction?. 2002 , 88, 137-41	29
685	Percutaneous Mechanical Reperfusion During Acute Myocardial Infarction. 2002 , 17, 162-173	O
684	Myocardial contrast echocardiography in acute coronary syndromes. 2002 , 17, 455-63	6
683	L-NAME enhances microcirculatory congestion and cardiomyocyte apoptosis during myocardial ischemia-reperfusion in rats. 2002 , 17, 185-92	14
682	Optimal time for predicting left ventricular remodeling after successful primary coronary angioplasty in acute myocardial infarction using serial myocardial contrast echocardiography and magnetic resonance imaging. 2002 , 66, 685-90	14
681	Impact of normalized myocardial perfusion after successful angioplasty in acute myocardial infarction. 2002 , 39, 591-7	322
68o	Repeated assessment of coronary flow velocity pattern in patients with first acute myocardial infarction. 2002 , 39, 1283-9	23
679	Effect of additional temporary glycoprotein IIb/IIIa receptor inhibition on troponin release in elective percutaneous coronary interventions after pretreatment with aspirin and clopidogrel (TOPSTAR trial). 2002 , 40, 662-8	96
678	Open perforator hypothesis: bridging epicardial and microvascular circulation. 2002, 40, 1214-5	
677	Combined assessment of microvascular integrity and contractile reserve improves differentiation of stunning and necrosis after acute anterior wall myocardial infarction. 2002 , 40, 1079-84	32
676	K(ATP) channel opening is an endogenous mechanism of protection against the no-reflow phenomenon but its function is compromised by hypercholesterolemia. 2002 , 40, 1339-46	53
675	Two different coronary blood flow velocity patterns in thrombolysis in myocardial infarction flow grade 2 in acute myocardial infarction: insight into mechanisms of microvascular dysfunction. 2002 , 40, 1755-60	30
674	Left ventricular electromechanical mapping: preliminary evidence of electromechanical changes after successful coronary intervention. 2002 , 144, 693-701	9
673	Embolization in atherosclerosis. 2002 , 12, 365-72	2
672	Intravenous myocardial contrast echocardiography predicts left ventricular remodeling in patients with acute myocardial infarction. 2002 , 15, 849-56	27
671	The significance of perfusion defect at myocardial perfusion MR imaging in a cat model of acute reperfused myocardial infarction. 2002 , 3, 235-9	1
,	with acute myocardial infarction. 2002 , 15, 849-56 The significance of perfusion defect at myocardial perfusion MR imaging in a cat model of acute	

670	Administration of glycoprotein IIb-IIIa inhibitors in patients with ST-segment elevation myocardial infarction. 2002 , 22, 864-88	2
669	Contrast echocardiography. 2002 , 27, 454-519	29
668	Reliability of resolution of ST-segment elevation after coronary reperfusion in predicting myocardial salvage in anterior wall acute myocardial infarction. 2002 , 90, 227-32	7
667	Frequency of incomplete reperfusion in patients with acute myocardial infarction undergoing primary angioplasty. 2002 , 90, 316-8	2
666	Usefulness of intravenous myocardial contrast echoardiography in the early left ventricular remodeling in acute myocardial infarction. 2002 , 90, 713-9	30
665	Myocardial contrast echocardiography in acute myocardial infarction. 2002 , 90, 38J-47J	22
664	Evaluation of myocardial viability with contrast echocardiography. 2002, 90, 65J-71J	28
663	Oxidative stress in septic shock and disseminated intravascular coagulation. 2002 , 33, 1173-85	142
662	Cardiac MRI: recent progress and continued challenges. 2002 , 16, 111-27	86
661	Percutaneous coronary intervention of thrombotic lesions: still challenging!. 2002 , 56, 8-9	6
660	Distal emboli protection in patients undergoing percutaneous coronary intervention after a recent myocardial infarction. 2002 , 57, 54-60	8
659	Intracoronary verapamil for reversal of no-reflow during coronary angioplasty for acute myocardial infarction. 2002 , 57, 444-51	86
658	The role of carnitine in myocardial dysfunction. 2003 , 41, S35-43	49
657	Effect of the PercuSurge GuardWire device on the integrity of microvasculature and clinical outcomes during primary transradial coronary intervention in acute myocardial infarction. 2003 , 92, 1331-5	94
656	Prognostic significance of transient no-reflow during primary percutaneous coronary intervention for ST-elevation acute myocardial infarction. 2003 , 92, 1445-7	49
655	Choc cardioghique primaire ^la phase aigulde lâlhfarctus trait par angioplastie : expfience de trois centres alsaciens publics non universitaires. 2003 , 52, 344-348	
654	Incremental value of myocardial contrast echocardiography for the prediction of recovery of function in dobutamine nonresponsive myocardium early after acute myocardial infarction. 2003 , 91, 397-402	46
653	ST-segment resolution and late (6-month) left ventricular remodeling after acute myocardial infarction. 2003 , 91, 451-3	13

(2003-2003)

652	Contrast echocardiographic evaluation of early changes in myocardial perfusion after recanalization therapy in anterior wall acute myocardial infarction and their relation with early contractile recovery. 2003 , 91, 532-7	11
651	Frequency and sequelae of ST elevation acute myocardial infarction caused by spontaneous distal embolization from unstable coronary lesions. 2003 , 91, 708-11	18
650	Inadvertent dilation of a saphenous vein graft stenosis by the PercuSurge GuardWire distal protection balloon. 2003 , 59, 346-9	5
649	Distal protection in native coronary arteries during primary angioplasty in acute myocardial infarction: single-center experience. 2003 , 60, 152-8	15
648	Evaluation of the PercuSurge Guardwire Plus Temporary Occlusion and Aspiration System during primary angioplasty in acute myocardial infarction. 2003 , 60, 443-51	21
647	The added clinical value of second generation ultrasound contrast agents. 2003 , 20 Suppl 1, S3-9	1
646	Assessment of myocardial viability in patients with postischemic left ventricular dysfunction: role of myocardial contrast echocardiography. 2003 , 20 Suppl 1, S19-29	2
645	Real-time perfusion imaging: a new echocardiographic technique for simultaneous evaluation of myocardial perfusion and contraction. 2003 , 20, 545-55	22
644	Contrast echocardiography using intravenous octafluoropropane and real-time perfusion imaging predicts functional recovery after acute myocardial infarction. 2003 , 16, 638-45	28
643	Clinical and angiographic correlates and outcomes of suboptimal coronary flow inpatients with acute myocardial infarction undergoing primary percutaneous coronary intervention. 2003 , 42, 1739-46	135
642	Association between hyperglycemia and the no-reflow phenomenon in patients with acute myocardial infarction. 2003 , 41, 1-7	276
641	Prediction of clinical outcome after mechanical revascularization in acute myocardial infarction by markers of myocardial reperfusion. 2003 , 41, 532-8	83
640	Detection of myocardial viability by contrast echocardiography in acute infarction predicts recovery of resting function and contractile reserve. 2003 , 41, 827-33	86
639	Thrombolysis in myocardial infarction myocardial perfusion grade in angiography correlates with myocardial salvage in patients with acute myocardial infarction treated with stenting or thrombolysis. 2003 , 41, 925-9	52
638	MR imaging of myocardial perfusion and viability. 2003 , 11, 49-66	28
637	Primary percutaneous transluminal coronary angioplasty accelerates early myocardial reperfusion compared to thrombolytic therapy in patients with acute myocardial infarction. 2003 , 146, 686-91	33
636	"Mirror-lake" serial relationship of electrocardiographic and biochemical indices for the detection of reperfusion and the prediction of salvage in patients with acute myocardial infarction. 2003 , 146, 757-63	7
635	Use of the electrocardiogram in acute myocardial infarction. 2003 , 348, 933-40	242

634	Clinical implications of ischaemia-reperfusion injury. 2003 , 9, 229-240	40
633	Hyperoxemic perfusion for treatment of reperfusion microvascular ischemia in patients with myocardial infarction. 2003 , 3, 253-63	22
632	Relation of coronary flow pattern to myocardial blush grade in patients with first acute myocardial infarction. 2003 , 89, 1147-51	30
631	Influence of contractile reserve and inducible ischaemia on left ventricular remodelling after acute myocardial infarction. 2003 , 89, 1138-43	19
630	Myocardial contrast echocardiography is superior to other known modalities for assessing myocardial reperfusion after acute myocardial infarction. 2003 , 89, 139-44	52
629	Impact of microvascular integrity and local viability on left ventricular remodelling after reperfused acute myocardial infarction. 2003 , 89, 393-7	26
628	Embolic protection devices. 2003 , 89, 990-2	28
627	Myocardial reperfusion: its assessment and its relation to clinical outcomes. 2003 , 1, 120-7	3
626	Effect of intracoronary estradiol on postischemic microvascular damage in a porcine model: a myocardial contrast echocardiographic study. 2003 , 54, 701-9	1
625	Persistence of systolic coronary flow reversal predicts irreversible dysfunction after reperfused anterior myocardial infarction. 2003 , 89, 382-8	18
625		18
	value of echocardiography in predicting future cardiac events after acute myocardial infarction.	
624	anterior myocardial infarction. 2003, 89, 382-8 Value of echocardiography in predicting future cardiac events after acute myocardial infarction. 2003, 18, 378-84 Impact of tirofiban on angiographic morphologic features of high-burden thrombus formation	3
624	anterior myocardial infarction. 2003, 89, 382-8 Value of echocardiography in predicting future cardiac events after acute myocardial infarction. 2003, 18, 378-84 Impact of tirofiban on angiographic morphologic features of high-burden thrombus formation during direct percutaneous coronary intervention and short-term outcomes. 2003, 124, 962-8 Facilitated primary percutaneous transluminal coronary angioplasty for acute ST segment elevation myocardial infarction: rationale for reuniting pharmacologic and mechanical revascularization	3 27
624 623	Value of echocardiography in predicting future cardiac events after acute myocardial infarction. 2003, 18, 378-84 Impact of tirofiban on angiographic morphologic features of high-burden thrombus formation during direct percutaneous coronary intervention and short-term outcomes. 2003, 124, 962-8 Facilitated primary percutaneous transluminal coronary angioplasty for acute ST segment elevation myocardial infarction: rationale for reuniting pharmacologic and mechanical revascularization strategies. 2003, 11, 13-20 Intravenous nicorandil in conjunction with coronary reperfusion therapy is associated with better	3 27 8
624 623 622	Value of echocardiography in predicting future cardiac events after acute myocardial infarction. 2003, 18, 378-84 Impact of tirofiban on angiographic morphologic features of high-burden thrombus formation during direct percutaneous coronary intervention and short-term outcomes. 2003, 124, 962-8 Facilitated primary percutaneous transluminal coronary angioplasty for acute ST segment elevation myocardial infarction: rationale for reuniting pharmacologic and mechanical revascularization strategies. 2003, 11, 13-20 Intravenous nicorandil in conjunction with coronary reperfusion therapy is associated with better clinical and functional outcomes in patients with acute myocardial infarction. 2003, 67, 295-300 Rescue percutaneous thrombectomy system provides better angiographic coronary flow and does	3 27 8 58
624 623 622 621	Value of echocardiography in predicting future cardiac events after acute myocardial infarction. 2003, 18, 378-84 Impact of tirofiban on angiographic morphologic features of high-burden thrombus formation during direct percutaneous coronary intervention and short-term outcomes. 2003, 124, 962-8 Facilitated primary percutaneous transluminal coronary angioplasty for acute ST segment elevation myocardial infarction: rationale for reuniting pharmacologic and mechanical revascularization strategies. 2003, 11, 13-20 Intravenous nicorandil in conjunction with coronary reperfusion therapy is associated with better clinical and functional outcomes in patients with acute myocardial infarction. 2003, 67, 295-300 Rescue percutaneous thrombectomy system provides better angiographic coronary flow and does not increase the in-hospital cost in patients with acute myocardial infarction. 2003, 67, 768-74 Relation between the timing of the last preinfarction angina and microvascular reperfusion in	3 27 8 58

616	[Early left ventricular remodelling following acute coronary accident]. 2004 , 20, 643-50	1
615	Contrast-enhanced MR imaging of the heart: overview of the literature. 2004 , 232, 653-68	119
614	Acute myocardial infarction: evaluation with first-pass enhancement and delayed enhancement MR imaging compared with 201Tl SPECT imaging. 2004 , 232, 49-57	166
613	The powerful microbubble: from bench to bedside, from intravascular indicator to therapeutic delivery system, and beyond. 2004 , 287, H450-7	165
612	Optimal therapeutic strategies in the setting of post-infarct no reflow: the need for a pathogenetic classification. 2004 , 90, 123-5	40
611	Myocardial Contrast Echocardiography in Acute Myocardial Infarction. 2004 , 191-218	
610	Clinical Application of Quantitative Analysis in Myocardial Contrast Echocardiography. 2004, 91-103	
609	The no-reflow phenomenon in coronary arteries. 2004 , 2, 1903-7	11
608	Serotonin blockade protects against early microvascular constriction following atherosclerotic plaque rupture. 2004 , 486, 85-9	4
607	Effects of percutaneous coronary arterial thrombectomy during acute myocardial infarction on left ventricular remodeling. 2004 , 93, 527-31	28
606	Prediction of the no-reflow phenomenon with ultrasonic tissue characterization in patients with anterior wall acute myocardial infarction. 2004 , 93, 1357-61, A5	7
605	Hemodynamics of microvascular dysfunction in patients with anterior wall acute myocardial infarction. 2004 , 94, 209-12	13
604	[Measurement of coronary flow and flow reserve with transthoracic echocardiography: an old concept, a new tool, a lot of applications]. 2004 , 53, 325-34	2
603	Impact of flow level on coronary flow velocity pattern. A doppler flow study in patients with first acute myocardial infarction. 2004 , 20, 27-35	O
602	Fourier phase analysis can be used to objectively analyze real-time myocardial contrast echocardiograms. 2004 , 20, 241-8	3
601	Relation of lateral ST-segment elevation pattern to myocardial salvage in patients with recanalized anterolateral acute myocardial infarction. 2004 , 27, 106-11	2
600	Thrombus aspiration prior to primary angioplasty in acute myocardial infarction: estimation of rescued myocardial tissue by return of ST-segment elevation. 2004 , 27, 451-4	2
599	Depletion of antioxidants is associated with no-reflow phenomenon in acute myocardial infarction. 2004 , 27, 466-70	19

Assessment of 99mTc-sestamibi myocardial redistribution following acute myocardial infarction and revascularization. **2004**, 24, 33-9

597	Angiographic predictors of left ventricular ejection fraction after successful angioplasty in acute myocardial infarction: an angiographic risk score for use in the catheterization laboratory. 2004 , 61, 338-43	3
596	Distal protection of bifurcating vessels: a novel approach. 2004 , 61, 512-4	2
595	Treatment of slow/no-reflow phenomenon with intracoronary nitroprusside injection in primary coronary intervention for acute myocardial infarction. 2004 , 63, 171-6	50
594	Improved performance of a new thrombus aspiration catheter: outcomes from in vitro experiments and a case presentation. 2004 , 63, 299-306	12
593	Contrast echocardiography: new agents. 2004 , 30, 425-34	63
592	Noninvasive assessment of coronary microvascular dysfunction using Tc-99m tetrofosmin SPECT in patients with acute myocardial infarction. 2004 , 11, 562-9	О
591	Acute pancreatitis mimicking myocardial infarction: evaluation with myocardial contrast echocardiography. 2004 , 17, 387-90	17
590	Myocardial contrast echocardiography predicts left ventricular remodelling after acute myocardial infarction. 2004 , 17, 1030-6	30
589	Myocardial contrast echocardiography in acute coronary syndromes. 2004 , 22, 253-67	2
588	Early assessment of coronary flow reserve by transthoracic Doppler echocardiography predicts late remodeling in reperfused anterior myocardial infarction. 2004 , 17, 750-5	39
587	Value of myocardial contrast echocardiography for predicting left ventricular remodeling and segmental functional recovery after anterior wall acute myocardial infarction. 2004 , 17, 923-32	20
586	Valor pron\(\text{Etico de marcadores no invasivos de reperfusi\(\text{E}\) coronaria frente a flujo TIMI 3 en pacientes tratados con angioplastia primaria. 2004 , 57, 524-530	1
585	Myocardial Contrast Echocardiography in Acute Myocardial Infarction: the Importance of Assessing Coronary Microcirculation. 2004 , 57, 4-6	О
584	Prognostic Value of Noninvasive Markers of Coronary Reperfusion Compared to TIMI 3 Flow in Patients Treated With Primary Angioplasty. 2004 , 57, 524-530	
583	Real Time Myocardial Contrast Echocardiography to Predict Left Ventricular Wall Motion Recovery After Reperfused Acute Myocardial Infarction. 2004 , 57, 815-825	О
582	Effects of glycoprotein IIb/IIIa inhibition on microvascular flow after coronary reperfusion. A quantitative myocardial contrast echocardiography study. 2004 , 43, 276-83	87
581	Enhanced coronary flow velocity during intra-aortic balloon pumping assessed by transthoracic Doppler echocardiography. 2004 , 43, 368-76	26

580	Glycoprotein IIb/IIIa inhibitors and no-reflow. 2004 , 43, 284-6	17
579	Reduction of "no-reflow" phenomenon by intra-aortic balloon counterpulsation in a randomized magnetic resonance imaging experimental study. 2004 , 43, 1291-8	47
578	Frequency, correlates, and clinical implications of myocardial perfusion after primary angioplasty and stenting, with and without glycoprotein IIb/IIIa inhibition, in acute myocardial infarction. 2004 , 44, 305-12	150
577	Evaluation of cardioprotective effects of recombinant soluble P-selectin glycoprotein ligand-immunoglobulin in myocardial ischemia-reperfusion injury by real-time myocardial contrast echocardiography. 2004 , 44, 887-91	22
576	Ecocardiografii de perfusiii miociidica en tiempo real para la predicciii de la recuperaciii de la funciii ventricular despus del infarto agudo de miocardio reperfundido. 2004 , 57, 815-825	O
575	Ecocardiografa de contraste en el infarto agudo de miocardio: la importancia de estudiar la microcirculacia coronaria. 2004 , 57, 4-6	
574	Detection of microvascular injury by evaluating epicardial blood flow in early reperfusion following primary angioplasty. 2004 , 96, 389-96	13
573	Myocardial endothelin-1 release and indices of inflammation during angioplasty for acute myocardial infarction and stable coronary artery disease. 2004 , 148, e10	17
572	Nicorandil improves cardiac function and clinical outcome in patients with acute myocardial infarction undergoing primary percutaneous coronary intervention: role of inhibitory effect on reactive oxygen species formation. 2004 , 148, E15	88
571	Infarction induced ventricular remodeling. 2004 , 78, 1507-10	2
570	Cardioprotective effects of magnesium sulfate in patients undergoing primary coronary angioplasty for acute myocardial infarction. 2004 , 68, 23-8	19
569	Incidence and management of "no-reflow" following percutaneous coronary interventions. 2005 , 329, 78-85	9
568	The effects of prior use of atorvastatin on coronary blood flow after primary percutaneous coronary intervention in patients presenting with acute myocardial infarction. 2005 , 16, 321-6	17
567	Automatic fuzzy classification of the washout curves from magnetic resonance first-pass perfusion imaging after myocardial infarction. 2005 , 40, 545-55	6
566	Plasminogen activator inhibitor-1 4G/5G polymorphism and efficacy of reperfusion therapy in acute myocardial infarction. 2005 , 16, 511-5	7
565	Evaluation of transmural myocardial perfusion by ultra-harmonic myocardial contrast echocardiography in reperfused acute myocardial infarction. 2005 , 69, 1041-6	7
564	Early administration of intracoronary verapamil improves myocardial perfusion during percutaneous coronary interventions for acute myocardial infarction. 2005 , 128, 2593-8	31
563	Efficacy of primary PCI: the microvessel perspective. 2005 , 7, I4-I9	3

562	The impact of admission C-reactive protein levels on the development of poor myocardial perfusion after primary percutaneous intervention in patients with acute myocardial infarction. 2005 , 16, 293-9	25
561	Persistent hyperglycemia is associated with left ventricular dysfunction in patients with acute myocardial infarction. 2005 , 69, 23-8	20
560	Reperfusion phenomenon is a strong predictor of left ventricular remodeling after acute myocardial infarction. 2005 , 69, 884-9	9
559	Intravenous versus intracoronary myocardial contrast echocardiography for evaluation of no-reflow after primary percutaneous coronary intervention. 2005 , 22, 818-25	2
558	Recovery of left ventricular systolic function after left anterior descending coronary artery stenting. 2005 , 18, 83-8	9
557	Assessment of myocardial viability using myocardial contrast echocardiography. 2005 , 22, 85-94	11
556	Fate of unprotected side branches as related to embolic complications during stent implantation for acute coronary syndromes using a distal protection procedure. 2005 , 95, 636-9	8
555	Usefulness of real-time myocardial perfusion imaging to evaluate tissue level reperfusion in patients with non-ST-elevation myocardial infarction. 2005 , 95, 1033-8	31
554	Effects of the nitric oxide donor nitroprusside on no-reflow phenomenon during coronary interventions for acute myocardial infarction. 2005 , 95, 1358-61	76
553	Usefulness of real-time intravenous myocardial contrast echocardiography in predicting left ventricular dilation after successfully reperfused acute myocardial infarction. 2005 , 96, 17-21	7
552	Usefulness of a new grading system based on coronary flow velocity pattern in predicting outcome in patients with acute myocardial infarction having percutaneous coronary intervention. 2005 , 96, 927-32	17
551	Prediction of decrease in myocardial perfusion defect size and severity during a 3-month follow-up by the degree of acute resolution of electrocardiographic changes. 2005 , 38, 100-5	4
550	Electrocardiogram of acute ST-elevation myocardial infarction: the significance of the various "scores". 2005 , 38, 113-8	3
549	Intracoronary ST segment evolution during primary coronary stenting predicts infarct zone recovery. 2005 , 64, 53-60	12
548	Brief repetitive balloon occlusions enhance reperfusion during percutaneous coronary intervention for acute myocardial infarction: a pilot study. 2005 , 65, 361-7	179
547	Coronary circulation and interventional cardiology. 2005 , 33, 1735-42	9
546	Cardiac imaging to identify patients at risk for developing heart failure after myocardial infarction. 2005 , 2, 183-8	14
545	Current Assessments of the Adequacy of Myocardial Perfusion During Acute MI. 2005 , 7, 25-34	

544	Ischemic Heart Disease. 2005 , 173-216	3
543	Predicting no-reflow based on angiographic features of lesions in patients with acute myocardial infarction. 2005 , 12, 315-21	8
542	Assessment of Myocardial Blood Volume. 2005 , 277-291	
541	Relation between the TIMI frame count and the degree of microvascular injury after primary coronary angioplasty in patients with acute anterior myocardial infarction. 2005 , 91, 64-7	14
540	Usefulness of myocardial parametric imaging to evaluate myocardial viability in experimental and in clinical studies. 2006 , 92, 350-6	4
539	The ECG in acute coronary syndromes: new tricks from an old dog. 2005 , 91, 851-3	7
538	Combined prognostic utility of ST-segment recovery and myocardial blush after primary percutaneous coronary intervention in acute myocardial infarction. 2005 , 26, 667-74	145
537	Sequelae of acute myocardial infarction regarding cardiac structure and function and their prognostic significance as assessed by magnetic resonance imaging. 2005 , 26, 549-57	400
536	Trends in myocardial contrast echocardiography: parameteric versus volumetric imaging. 2005, 3, 133-42	1
535	Intravenous myocardial contrast echocardiography predicts regional and global left ventricular remodelling after acute myocardial infarction: comparison with low dose dobutamine stress echocardiography. 2005 , 91, 1578-83	15
534	The effect of interventional treatment in acute myocardial infarction on ST resolution: a comparison of coronary angioplasty with excimer laser angioplasty. 2005 , 56, 377-84	10
533	The effect of intra-aortic balloon counterpulsation on left ventricular functional recovery early after acute myocardial infarction: a randomized experimental magnetic resonance imaging study. 2005 , 26, 1235-41	34
532	Infarct zone viability influences ventricular remodelling after late recanalisation of an occluded infarct related artery. 2005 , 91, 478-83	28
531	Tratamiento farmacol?gico del infarto agudo de miocardio. Fibrin?lisis. Criterios de respuesta y seguimiento. 2005 , 9, 2653-2661	
530	Myocardial ischemia/reperfusion-injury, a clinical view on a complex pathophysiological process. 2005 , 100, 179-90	339
529	Potential clinical applications of myocardial contrast echocardiography in evaluating myocardial perfusion in coronary artery disease. 2005 , 104, 1-9	5
528	Very early cardiac magnetic resonance imaging for quantification of myocardial tissue perfusion in patients receiving tirofiban before percutaneous coronary intervention for ST-elevation myocardial infarction. 2005 , 149, 564	27
527	ST resolution in a single electrocardiographic lead: a simple and accurate predictor of cardiac mortality in patients with fibrinolytic therapy for acute ST-elevation myocardial infarction. 2005 , 149, 91-7	36

526	Acute hyperglycemia is associated with adverse outcome after acute myocardial infarction in the coronary intervention era. 2005 , 150, 814-20	138
525	Distal embolization during primary angioplasty: histopathologic features and predictability. 2005 , 150, 102-8	80
524	Left ventricular remodeling after primary coronary angioplasty in patients treated with abciximab or intracoronary adenosine. 2005 , 150, 1015	60
523	Mean platelet volume on admission predicts impaired reperfusion and long-term mortality in acute myocardial infarction treated with primary percutaneous coronary intervention. 2005 , 46, 284-90	266
522	Leucocyte-endothelial interactions in health and disease. 2006 , 97-133	64
521	Association of C-reactive protein and myocardial perfusion in patients with ST-elevation acute myocardial infarction. 2006 , 186, 177-83	26
520	Coronary flow velocity pattern assessed noninvasively by transthoracic color Doppler echocardiography serves as a predictor of adverse cardiac events and left ventricular remodeling in patients with acute myocardial infarction. 2006 , 19, 335-40	13
519	Automated quantification of the spatial extent of perfusion defects and viability on myocardial contrast echocardiography. 2006 , 19, 379-85	7
518	Value of Coronary Blood Flow Pattern as a Predictor of Functional Recovery and Short-Term Left Ventricular Remodeling After Primary Coronary Angioplasty. A Transthoracic Doppler Study. 2006 , 59, 352-359	
517	Coronary Angiography: Beyond Coronary Anatomy. 2006 , 59, 596-608	1
516	Valor del patrfi de flujo coronario tras angioplastia primaria como predictor de recuperacifi funcional y remodelado ventricular a corto plazo. Estudio mediante ecocardiograffi Doppler transtorfica. 2006 , 59, 352-359	O
515	Coronariografā: mā allīde la anatomā coronaria. 2006 , 59, 596-608	3
514	No-reflow phenomenon and prognosis in patients with acute myocardial infarction. 2006 , 3, 499-506	144
513	Final myocardial blush grade predicts troponin I elevation in unstable angina patients undergoing percutaneous intervention. 2006 , 15, 291-6	
512	Sestamibi single photon emission computed tomography immediately after primary percutaneous coronary intervention identifies patients at risk for large infarcts. 2006 , 151, 1108-14	13
512 511		13 32
	coronary intervention identifies patients at risk for large infarcts. 2006 , 151, 1108-14 Thrombus Aspiration during Percutaneous coronary intervention in Acute myocardial infarction	

(2006-2006)

508	Nicorandil reduces myocardial no-reflow by protection of endothelial function via the activation of KATP channel. 2006 , 374, 100-5		18
507	Evaluation of no-reflow phenomenon using 201TlCl/123I-BMIPP dual-isotope myocardial SPECT. 2006 , 73, 258-64		3
506	The Importance of Amino-terminal pro-Brain Natriuretic Peptide Testing in Clinical Cardiology. <i>Biomarker Insights</i> , 2006 , 1, 117727190600100	3.5	О
505	Continuous administration of nicorandil decreases QT dispersion during the chronic phase of acute myocardial infarction. 2006 , 47, 351-61		16
504	Impact of nicorandil to prevent reperfusion injury in patients with acute myocardial infarction: Sigmart Multicenter Angioplasty Revascularization Trial (SMART). 2006 , 70, 1099-104		61
503	Effect of distal protection device on the microvascular integrity in acute myocardial infarction during primary percutaneous coronary intervention. 2006 , 70, 1284-9		5
502	Impact of PercuSurge device conjugative with intracoronary administration of nitroprusside on no-reflow phenomenon following primary percutaneous coronary intervention. 2006 , 70, 1538-42		18
501	Pressure-derived collateral flow index: a strong predictor of late left ventricular remodeling after thrombolysis for acute myocardial infarction. 2006 , 17, 139-44		9
500	Pretreatment with fosinopril or valsartan reduces myocardial no-reflow after acute myocardial infarction and reperfusion. 2006 , 17, 463-9		21
499	Coronary flow velocity pattern assessed by transthoracic Doppler echocardiography predicts adverse clinical events and myocardial recovery after successful primary angioplasty. 2006 , 7, 753-60		3
498	Contrast echocardiography for the assessment of myocardial viability. 2006 , 21, 473-8		18
497	Effects of intraaortic balloon pumping on the angiographic no-reflow phenomenon after percutaneous coronary intervention in patients with anterior myocardial infarction. 2006 , 70, 37-43		12
496	Value of adenosine infusion for infarct size determination using real-time myocardial contrast echocardiography. 2006 , 4, 10		1
495	Usefulness of myocardial contrast echocardiography early after acute myocardial infarction. 2006 , 23, 208-17		3
494	Pretreatment with simvastatin reduces myocardial no-reflow by opening mitochondrial K(ATP) channel. 2006 , 149, 243-9		38
493	Imaging microvascular obstruction and its clinical significance following acute myocardial infarction. 2006 , 11, 305-12		19
492	Assessment of resting perfusion defects in patients with acute myocardial infarction: comparison of myocardial contrast echocardiography, combined first-pass/delayed contrast-enhanced magnetic resonance imaging and 99mTC-sestamibi SPECT. 2006 , 22, 417-28		13
491	Different effects of adenosine and calcium channel blockade on myocardial no-reflow after acute myocardial infarction and reperfusion. <i>Cardiovascular Drugs and Therapy</i> , 2006 , 20, 167-75	3.9	6

490	Assessment of no-reflow regions using cardiac MRI. 2006 , 101, 383-90	51
489	Evaluating the 'no reflow' phenomenon with myocardial contrast echocardiography. 2006 , 101, 391-9	16
488	Combination study of myocardial perfusion and left ventricular filling provides an excellent prediction of clinical outcomes in patients with reperfused myocardial infarction. 2006 , 101, 400-7	9
487	Hyperbaric oxygen therapy prevents coagulation disorders in an experimental model of multiple organ failure syndrome. 2006 , 32, 1881-8	13
486	Comparison between contrast echocardiography and magnetic resonance imaging to predict improvement of myocardial function after primary coronary intervention. 2006 , 97, 361-6	17
485	Influence of transmurality, infarct size, and severe microvascular obstruction on left ventricular remodeling and function after primary coronary angioplasty. 2006 , 98, 1033-40	114
484	ST-segment recovery and prognosis in patients with ST-elevation myocardial infarction reperfused by prehospital combination fibrinolysis, prehospital initiated facilitated percutaneous coronary intervention, or primary percutaneous coronary intervention. 2006 , 98, 1132-9	27
483	Usefulness of myocardial contrast echocardiography in predicting late mortality in patients with anterior wall acute myocardial infarction. 2006 , 98, 1150-5	26
482	Contrast-enhanced cardiac magnetic resonance in the evaluation of myocardial infarction and myocardial viability in patients with ischemic heart disease. 2006 , 31, 128-68	23
481	Modern management of acute myocardial infarction. 2006 , 31, 769-817	11
481 480	Modern management of acute myocardial infarction. 2006, 31, 769-817 Tratamiento coadyuvante en el intervencionismo coronario percutileo del infarto agudo de miocardio. 2006, 6, 39H-48H	11
	Tratamiento coadyuvante en el intervencionismo coronario percutfieo del infarto agudo de	11
480	Tratamiento coadyuvante en el intervencionismo coronario percutibeo del infarto agudo de miocardio. 2006 , 6, 39H-48H	17
480 479	Tratamiento coadyuvante en el intervencionismo coronario percutileo del infarto agudo de miocardio. 2006, 6, 39H-48H Valoracifi de la viabilidad miocidica mediante resonancia magnica. 2006, 6, 49E-56E Hyperoxemic perfusion of the left anterior descending coronary artery after primary angioplasty in	
480 479 478	Tratamiento coadyuvante en el intervencionismo coronario percutfieo del infarto agudo de miocardio. 2006, 6, 39H-48H Valoracifi de la viabilidad miocfidica mediante resonancia magnica. 2006, 6, 49E-56E Hyperoxemic perfusion of the left anterior descending coronary artery after primary angioplasty in anterior ST-elevation myocardial infarction. 2006, 67, 859-65 Potential significance of spontaneous and interventional ST-changes in patients transferred for primary percutaneous coronary intervention: observations from the ST-MONitoring in Acute	17
480 479 478 477	Tratamiento coadyuvante en el intervencionismo coronario percutileo del infarto agudo de miocardio. 2006, 6, 39H-48H Valoracifi de la viabilidad miocidica mediante resonancia magnitica. 2006, 6, 49E-56E Hyperoxemic perfusion of the left anterior descending coronary artery after primary angioplasty in anterior ST-elevation myocardial infarction. 2006, 67, 859-65 Potential significance of spontaneous and interventional ST-changes in patients transferred for primary percutaneous coronary intervention: observations from the ST-MONitoring in Acute Myocardial Infarction study (The MONAMI study). 2006, 27, 267-75 Microvascular obstruction and myocardial function after acute myocardial infarction: assessment	17 57
480 479 478 477 476	Tratamiento coadyuvante en el intervencionismo coronario percutileo del infarto agudo de miocardio. 2006, 6, 39H-48H Valoracifi de la viabilidad miocidica mediante resonancia magnitica. 2006, 6, 49E-56E Hyperoxemic perfusion of the left anterior descending coronary artery after primary angioplasty in anterior ST-elevation myocardial infarction. 2006, 67, 859-65 Potential significance of spontaneous and interventional ST-changes in patients transferred for primary percutaneous coronary intervention: observations from the ST-MONitoring in Acute Myocardial Infarction study (The MONAMI study). 2006, 27, 267-75 Microvascular obstruction and myocardial function after acute myocardial infarction: assessment by using contrast-enhanced cine MR imaging. 2006, 240, 529-36 Impact of intracoronary aspiration thrombectomy during primary angioplasty on left ventricular	17 57 17

(2007-2007)

472	Troponin-I concentration 72 h after myocardial infarction correlates with infarct size and presence of microvascular obstruction. 2007 , 93, 1547-51	56
471	Distal embolic protection during percutaneous coronary intervention in patients with acute coronary syndromes: the RUBY study. 2006 , 8, 148-54	8
470	Different effects of tirofiban and aspirin plus clopidogrel on myocardial no-reflow in a mini-swine model of acute myocardial infarction and reperfusion. 2006 , 92, 1131-7	35
469	The open artery hypothesis: beneficial effects and long-term prognostic importance of patency of the infarct-related coronary artery. 2007 , 58, 34-44	13
468	Remote ischaemic postconditioning protects the heart during acute myocardial infarction in pigs. 2007 , 93, 749-52	147
467	Post-infarction treatment with simvastatin reduces myocardial no-reflow by opening of the KATP channel. 2007 , 9, 30-6	21
466	Impaired perfusion after myocardial infarction is due to reperfusion-induced deltaPKC-mediated myocardial damage. 2007 , 73, 699-709	40
465	Prediction of left ventricular remodeling and analysis of infarct resorption in patients with reperfused myocardial infarcts by using contrast-enhanced MR imaging. 2007 , 245, 95-102	84
464	Percutaneous coronary intervention and the no-reflow phenomenon. 2007 , 5, 715-31	27
463	Reperfusion arrhythmias: are they only a marker of epicardial reperfusion or continuing myocardial ischemia after acute myocardial infarction?. 2007 , 58, 663-70	13
462	Myocardial blush grade: to evaluate myocardial viability in patients with acute myocardial infarction. 2007 , 58, 556-60	1
461	Atherosclerotic plaque with ultrasonic attenuation affects coronary reflow and infarct size in patients with acute coronary syndrome: an intravascular ultrasound study. 2007 , 71, 648-53	70
460	Intravenous contrast echocardiography after myocardial infarction: relationship among residual myocardial perfusion, contractile reserve and long-term remodelling. 2007 , 8, 1012-9	3
459	Understanding the heart: CT and MRI for coronary heart disease. 2007 , 22, 107-13	5
458	Primary angioplasty and routine utilization of thrombus aspiration devices: feasibility and results in a consecutive series of 486 patients. 2007 , 8, 258-64	11
457	Quantitative evaluation of myocardial blush to assess tissue level reperfusion in patients with acute ST-elevation myocardial infarction: incremental prognostic value compared with visual assessment. 2007 , 153, 612-20	28
456	Different effects of postconditioning on myocardial no-reflow in the normal and hypercholesterolemic mini-swines. 2007 , 73, 137-42	47
455	Recovery of global systolic function after primary angioplasty. Influence of coronary flow velocity reserve measured by transthoracic echocardiography. 2007 , 114, 315-22	2

454	Effect of microvascular reperfusion on prognosis and left ventricular function in anterior wall myocardial infarction treated with primary angioplasty. 2007 , 114, 183-7	13
453	Carvedilol preserves endothelial junctions and reduces myocardial no-reflow after acute myocardial infarction and reperfusion. 2007 , 115, 334-41	15
452	Excimer laser thrombus elimination for prevention of distal embolization and no-reflow in patients with acute ST elevation myocardial infarction: results from the randomized LaserAMI study. 2007 , 116, 20-6	40
451	An increased platelet-leukocytes interaction at the culprit site of coronary artery occlusion in acute myocardial infarction: a pathogenic role for "no-reflow" phenomenon?. 2007 , 117, 123-30	44
450	Role of pre-infarction angina and inflammatory status in the extent of microvascular obstruction detected by MRI in myocardial infarction patients treated by PCI. 2007 , 121, 139-47	27
449	Coronary microvascular dysfunction. 2007 , 356, 830-40	1151
448	Stem cell therapy improves myocardial perfusion and cardiac synchronicity: new application for echocardiography. 2007 , 20, 512-20	11
447	Significance of additional ST segment elevation in patients with no reflow after angioplasty for acute myocardial infarction. 2007 , 20, 262-9	1
446	Retinal arteriolar narrowing and left ventricular remodeling: the multi-ethnic study of atherosclerosis. 2007 , 50, 48-55	114
445	Prognostic value of myocardial viability detected by myocardial contrast echocardiography early after acute myocardial infarction. 2007 , 50, 327-34	65
444	The Usefulness of Intracoronary Electrocardiography during Primary Percutaneous Coronary Intervention in Patients with Acute Myocardial Infarction. <i>Korean Circulation Journal</i> , 2007 , 37, 148	
443	Selective intracoronary administration of nitroprusside before balloon dilatation prevents slow reflow during percutaneous coronary intervention in patients with acute myocardial infarction. 2007 , 48, 423-33	28
442	Relation of hyperglycemia to ST-segment resolution after primary percutaneous coronary intervention for acute myocardial infarction. 2007 , 120, 1874-1877	1
441	Bone-marrow-derived Cells in Myocardial Repair and Regeneration. 2007, 117-127	
440	High dose adenosine for suboptimal myocardial reperfusion after primary PCI: A randomized placebo-controlled pilot study. 2008 , 71, 283-9	61
439	MRI evaluation of microvascular obstruction in experimental reperfused acute myocardial infarction using a T1 and T2 preparation pulse sequence. 2007 , 26, 1486-92	7
438	Proximal coronary hemodynamic changes evaluated by intracardiac echocardiography during myocardial ischemia and reperfusion in a canine model. 2008 , 25, 312-20	2
437	Prospective, multicenter study of thrombectomy in patients with acute myocardial infarction: the X-Tract AMI registry. 2007 , 20, 44-50	14

(2008-2007)

436	myocardial infarction. 2007 , 20, 340-50	10
435	Contrast echocardiography. 2007 , 32, 51-96	104
434	Frequency of slow coronary flow following successful stent implantation and effect of Nitroprusside. 2007 , 99, 916-20	25
433	Usefulness of coronary flow reserve immediately after primary coronary angioplasty for acute myocardial infarction in predicting long-term adverse cardiac events. 2007 , 100, 806-11	22
432	Prediction of improvement in left ventricular function during a 1-year follow-up after acute myocardial infarction by the degree of acute resolution of electrocardiographic changes. 2007 , 40, 416-21	3
431	Chronic pretreatment of ACEI reduces no-reflow in patients with acute myocardial infarction treated with primary angioplasty. 2007 , 30, 130-4	9
430	Determinants and impact of microvascular obstruction in successfully reperfused ST-segment elevation myocardial infarction. Assessment by magnetic resonance imaging. 2007 , 17, 2572-80	106
429	Intracoronary boluses of adenosine and sodium nitroprusside in combination reverses slow/no-reflow during angioplasty: a clinical scenario of ischemic preconditioning. 2007 , 85, 476-82	29
428	Assessment of non-reperfused and reperfused myocardial infarction using diffusible or deposited radiolabelled perfusion imaging agents. 2007 , 34, 330-7	1
427	Upfront thrombus aspiration in primary coronary intervention for patients with ST-segment elevation acute myocardial infarction: report of the VAMPIRE (VAcuuM asPIration thrombus REmoval) trial. 2008 , 1, 424-31	112
426	Evaluation of the left ventricular remodeling in patients with myocardial infarction after revascularization with intravenous real-time myocardial contrast echocardiography. 2008 , 28, 287-90	2
425	Myocardial perfusion imaging with contrast echocardiography. 2008 , 10, 233-9	13
424	Prognostic value of ST-segment resolution after rescue percutaneous coronary intervention. Data from the RICO survey. 2008 , 71, 607-12	5
423	Left ventricular remodeling can be predicted with left ventricular volume response during dobutamine echocardiography after acute myocardial infarction. 2008 , 31, 259-64	5
422	Evaluation cardioprotective effects of atorvastatin in rats by real time myocardial contrast echocardiography. 2008 , 25, 974-81	8
421	Effect of statin therapy before Q-wave myocardial infarction on myocardial perfusion. 2008, 101, 139-43	17
420	Assessment and key targets for therapy in the post-myocardial infarction patient with left ventricular dysfunction. 2008 , 102, 5G-12G	4
419	The post-myocardial infarction scarred ventricle and congestive heart failure: the preeminence of magnetic resonance imaging for preoperative, intraoperative, and postoperative assessment. 2008 , 136, 1405-12	29

418	Relation of admission QRS duration with development of angiographic no-reflow in patients with acute ST-segment elevation myocardial infarction treated with primary percutaneous interventions. 2008 , 41, 72-7	8
417	Myocardial ischemia: current concepts and future perspectives. 2008, 52, 67-78	72
416	Advances in the assessment of no-reflow after successful primary percutaneous coronary intervention for acute ST-segment elevation myocardial infarction: now that we can diagnose it, what do we do about it?. 2008 , 51, 566-8	10
415	Functional recovery after acute myocardial infarction: comparison between angiography, electrocardiography, and cardiovascular magnetic resonance measures of microvascular injury. 2008 , 52, 181-9	279
414	New echocardiographic techniques for evaluating left ventricular myocardial function. 2008, 12, 228-47	22
413	Angiographic and physiologic assessment of coronary flow and myocardial perfusion in the cardiac catheterization laboratory. 2008 , 10, 69-78	3
412	Effect of systemic blood pressure on microcollateral circulation evaluated by real-time contrast echocardiography. 2008 , 21, 765-9	8
411	Minimally invasive video-assisted thyroidectomy: the learning curve. 2008 , 41, 33-6	37
410	Myocyte Protection by Device Therapy. Fundamental and Clinical Cardiology, 2008, 248-264	
409	Contrast echocardiography: evidence-based recommendations by European Association of Echocardiography. 2009 , 10, 194-212	233
408	Randomized comparison between tirofiban and abciximab to promote complete ST-resolution in primary angioplasty: results of the facilitated angioplasty with tirofiban or abciximab (FATA) in ST-elevation myocardial infarction trial. 2008 , 29, 2972-80	35
408 407	primary angioplasty: results of the facilitated angioplasty with tirofiban or abciximab (FATA) in	35 1774
·	primary angioplasty: results of the facilitated angioplasty with tirofiban or abciximab (FATA) in ST-elevation myocardial infarction trial. 2008, 29, 2972-80 Management of acute myocardial infarction in patients presenting with persistent ST-segment elevation: the Task Force on the Management of ST-Segment Elevation Acute Myocardial Infarction	
407	primary angioplasty: results of the facilitated angioplasty with tirofiban or abciximab (FATA) in ST-elevation myocardial infarction trial. 2008, 29, 2972-80 Management of acute myocardial infarction in patients presenting with persistent ST-segment elevation: the Task Force on the Management of ST-Segment Elevation Acute Myocardial Infarction of the European Society of Cardiology. 2008, 29, 2909-45 Intravenous administration of nicorandil immediately before percutaneous coronary intervention	1774
407 406	primary angioplasty: results of the facilitated angioplasty with tirofiban or abciximab (FATA) in ST-elevation myocardial infarction trial. 2008, 29, 2972-80 Management of acute myocardial infarction in patients presenting with persistent ST-segment elevation: the Task Force on the Management of ST-Segment Elevation Acute Myocardial Infarction of the European Society of Cardiology. 2008, 29, 2909-45 Intravenous administration of nicorandil immediately before percutaneous coronary intervention can prevent slow coronary flow phenomenon. 2009, 30, 765-72 Beneficial effect of post-procedural abciximab in patients undergoing primary coronary angioplasty	1774 39
407 406 405	primary angioplasty: results of the facilitated angioplasty with tirofiban or abciximab (FATA) in ST-elevation myocardial infarction trial. 2008, 29, 2972-80 Management of acute myocardial infarction in patients presenting with persistent ST-segment elevation: the Task Force on the Management of ST-Segment Elevation Acute Myocardial Infarction of the European Society of Cardiology. 2008, 29, 2909-45 Intravenous administration of nicorandil immediately before percutaneous coronary intervention can prevent slow coronary flow phenomenon. 2009, 30, 765-72 Beneficial effect of post-procedural abciximab in patients undergoing primary coronary angioplasty and presenting with the no-reflow phenomenon. 2008, 10, 100-3 Role of adjunctive thrombectomy and embolic protection devices in acute myocardial infarction: a	1774 39 2
407 406 405 404	primary angioplasty: results of the facilitated angioplasty with tirofiban or abciximab (FATA) in ST-elevation myocardial infarction trial. 2008, 29, 2972-80 Management of acute myocardial infarction in patients presenting with persistent ST-segment elevation: the Task Force on the Management of ST-Segment Elevation Acute Myocardial Infarction of the European Society of Cardiology. 2008, 29, 2909-45 Intravenous administration of nicorandil immediately before percutaneous coronary intervention can prevent slow coronary flow phenomenon. 2009, 30, 765-72 Beneficial effect of post-procedural abciximab in patients undergoing primary coronary angioplasty and presenting with the no-reflow phenomenon. 2008, 10, 100-3 Role of adjunctive thrombectomy and embolic protection devices in acute myocardial infarction: a comprehensive meta-analysis of randomized trials. 2008, 29, 2989-3001 Incidence and clinical consequences of distal embolization on the coronary angiogram after	1774 39 2 193

(2009-2008)

400	in patients with acute ST segment elevation myocardial infarction (STEMI) treated with fibrinolytic therapy. 2008 , 72, 873-9	7
399	Relationship of admission neutrophil count to microvascular injury, left ventricular dilation, and long-term outcome in patients treated with primary angioplasty for acute myocardial infarction. 2008 , 72, 867-72	31
398	Predictors and long-term prognostic implications of angiographic slow/no-flow during percutaneous coronary intervention for acute myocardial infarction. 2008 , 47, 899-906	11
397	Evaluation of myocardial perfusion and deformation in patients with acute myocardial infarction treated with primary angioplasty and stent placement. 2008 , 19, 497-506	9
396	Impact of admission glomerular filtration rate on the development of poor myocardial perfusion after primary percutaneous intervention in patients with acute myocardial infarction. 2008 , 19, 543-9	7
395	Using a contrast-enhanced imaging sequence at 3-minute delay in 3-T magnetic resonance imaging for acute infarct evaluation. 2008 , 43, 669-75	11
394	Evaluation of acute coronary syndromes by cardiac magnetic resonance imaging. 2008, 19, 25-32	10
393	Ischemia-Reperfusion Injury and Hyperbaric Oxygen Therapy. 2008 , 159-185	3
392	Dose-dependent effects of intravenous allogeneic mesenchymal stem cells in the infarcted porcine heart. 2009 , 18, 321-9	32
391	Lipid-rich plaque and myocardial perfusion after successful stenting in patients with non-ST-segment elevation acute coronary syndrome: an optical coherence tomography study. 2009 , 30, 1348-55	151
390	Outcome after primary percutaneous intervention in acute myocardial infarction: role of microcirculatory perfusion a crucial piece in the puzzle. 2009 , 30, 530-1	2
389	Assessment of microvascular obstruction and prediction of short-term remodeling after acute myocardial infarction: cardiac MR imaging study. 2009 , 250, 363-70	98
388	Role of thrombectomy in acute myocardial infarction. 2009 , 7, 289-97	
387	The utilization of ultrasound and microbubbles for therapy in acute coronary syndromes. 2009 , 83, 636-42	14
386	Diagnostic ultrasound combined with glycoprotein IIb/IIIa-targeted microbubbles improves microvascular recovery after acute coronary thrombotic occlusions. <i>Circulation</i> , 2009 , 119, 1378-85	88
385	ST changes before and during primary percutaneous coronary intervention predict final infarct size in patients with ST elevation myocardial infarction. 2009 , 42, 64-72	22
384	Effect of intracoronary thrombectomy on 30-day mortality in non-diabetic patients with acute hyperglycemia after acute myocardial infarction. 2009 , 53, 429-36	5
383	Predictors of no-reflow after percutaneous coronary intervention for culprit lesion with plaque rupture in infarct-related artery in patients with acute myocardial infarction. 2009 , 54, 36-44	23

382	Comparison between myocardial contrast echocardiography and (99m)technetium sestamibi single photon emission computed tomography determined myocardial viability in predicting hard cardiac events following acute myocardial infarction. 2009 , 104, 1184-8	10
381	Myocardium selective densitometric perfusion assessment after acute myocardial infarction. 2009 , 10, 49-54	5
380	Comparison of the angiographic myocardial blush grade with delayed-enhanced cardiac magnetic resonance for the assessment of microvascular obstruction in acute myocardial infarctions. 2009 , 74, 1000-7	29
379	Participation of mast cells in angiogenesis in the border zone of myocardial infarction in rats. 2009 , 36, 119-27	2
378	Association of angiographic perfusion score following percutaneous coronary intervention for ST-elevation myocardial infarction with left ventricular remodeling at 6 weeks in GRACIA-2. 2009 , 27, 253-8	4
377	Factors associated with myocardial salvage immediately after emergent percutaneous coronary intervention in patients with ST-elevation acute myocardial infarction. 2009 , 23, 383-90	2
376	Scintigraphic evaluation of routine filterwire distal protection in percutaneous coronary intervention for acute ST-segment elevation myocardial infarction: a randomized controlled trial. 2009 , 16, 784-91	10
375	Non-invasive imaging of microvascular damage. 2009 , 16, 811-31	4
374	Appearance of microvascular obstruction on high resolution first-pass perfusion, early and late gadolinium enhancement CMR in patients with acute myocardial infarction. 2009 , 11, 33	71
373	Combined assessment of myocardial perfusion and diastolic function enhances risk stratification in patients with anterior wall myocardial infarction. 2009 , 26, 61-5	2
372	Clinical significance of post-procedural TIMI flow in patients with cardiogenic shock undergoing primary percutaneous coronary intervention. 2009 , 2, 56-64	41
371	Major prognostic impact of persistent microvascular obstruction as assessed by contrast-enhanced cardiac magnetic resonance in reperfused acute myocardial infarction. 2009 , 19, 2117-26	61
370	Remote periconditioning reduces myocardial no-reflow by the activation of K ATP channel via inhibition of Rho-kinase. 2009 , 133, 179-84	17
369	The impact of admission C-reactive protein levels on the development of no-reflow phenomenon after primary PCI in patients with acute myocardial infarction: the role of inflammation. 2009 , 136, 86-8; author reply 88-9	17
368	Role of inflammation in the extent of microvascular obstruction in patients undergoing primary PCI. 2009 , 135, 273-5	3
367	Cardiovascular magnetic resonance in patients with myocardial infarction: current and emerging applications. 2009 , 55, 1-16	250
366	Early electrocardiographic findings and MR imaging-verified microvascular injury and myocardial infarct size. 2009 , 2, 1187-94	39
365	The effect of statins on the no-reflow phenomenon: an observational study in patients with hyperglycemia before primary angioplasty. 2009 , 9, 81-9	15

(2010-2009)

364	Relationship of admission hematological indexes with myocardial reperfusion abnormalities in acute ST segment elevation myocardial infarction patients treated with primary percutaneous coronary interventions. 2009 , 25, e164-8	25
363	[Not Available]. 2009 , 62, 293.e1-293.e47	5
362	Contrast Echocardiography in Acute Coronary Syndromes. 2009 , 117-131	
361	End-organ dysfunction and cardiovascular outcomes: the role of the microcirculation. 2009 , 116, 175-90	49
360	Acute myocardial infarction: the enduring challenge for cardiac protection and survival. 2009 , 73, 2000-8	53
359	Early extracorporeal membrane oxygenator-assisted primary percutaneous coronary intervention improved 30-day clinical outcomes in patients with ST-segment elevation myocardial infarction complicated with profound cardiogenic shock. 2010 , 38, 1810-7	270
358	Limitations and potential clinical application on contrast echocardiography. 2010 , 6, 24-30	6
357	Non-invasive coronary flow reserve after successful primary angioplasty for acute anterior myocardial infarction is an independent predictor of left ventricular adverse remodelling. 2010 , 11, 711-8	11
356	Syndromes coronaires aigus. 2010 , 5, 1-21	
355	Predictive value of ischemic mitral regurgitation during the acute phase of ST elevation myocardial infarction treated with primary coronary intervention for left ventricular remodeling in long-term follow-up. 2010 , 21, 325-9	6
354	Post-reperfusion enhancement of CD14(+)CD16(-) monocytes and microvascular obstruction in ST-segment elevation acute myocardial infarction. 2010 , 74, 1175-82	19
353	High index of microcirculatory resistance level after successful primary percutaneous coronary intervention can be improved by intracoronary administration of nicorandil. 2010 , 74, 909-15	23
352	Rheolytic thrombectomy: any role left?. 2010 , 2, 57-65	2
351	Predictors and long-term prognosis of angiographic slow/no-reflow phenomenon during emergency percutaneous coronary intervention for ST-elevated acute myocardial infarction. 2010 , 33, E7-12	49
350	Injectable materials for the treatment of myocardial infarction and heart failure: the promise of decellularized matrices. 2010 , 3, 478-86	140
349	Hyperglykthie. 2010 , 4, 488-496	
348	How to mediate cardioprotection in ischemic heartsaccumulated evidence of basic research should translate to clinical medicine. <i>Cardiovascular Drugs and Therapy</i> , 2010 , 24, 217-23	7
347	Efficacy of thrombectomy for acute myocardial infarction-Special focus on its efficacy according to different infarct-related arteries. 2010 , 55, 189-95	6

346	Therapeutic advances in myocardial microvascular resistance: unravelling the enigma. 2010 , 127, 131-47	23
345	Impact of thrombus aspiration on angiographic and clinical outcomes in patients with ST-elevation myocardial infarction. 2010 , 11, 218-22	5
344	Dipyridamole versus verapamil for treatment of no-reflow during primary angioplasty. 2010 , 76, 787-93	10
343	Long-term outcome after thrombectomy in acute myocardial infarction. 2010 , 40, 233-41	6
342	Pathophysiology of myocardial injury and remodeling: implications for molecular imaging. 2010 , 51 Suppl 1, 102S-106S	12
341	Stenting of complex lesions: an overview. 2010 , 7, 485-96	37
340	Predicting the no-reflow phenomenon following successful percutaneous coronary intervention. 2010 , 4, 403-20	5
339	Les cellules màullaires traitès par onde de choc amliorent la fonction ventriculaire gauche aprà infarctus du myocarde chez le lapin. 2010 , 24, 882-895	
338	Serum soluble ST2: a potential novel mediator in left ventricular and infarct remodeling after acute myocardial infarction. 2010 , 55, 243-50	216
337	Microvascular obstruction: underlying pathophysiology and clinical diagnosis. 2010 , 55, 1649-60	
331	Microvascular obstruction: underlying pachophysiology and clinical diagnosis. 2010, 33, 1047 00	191
336	Value of a new multiparametric score for prediction of microvascular obstruction lesions in ST-segment elevation myocardial infarction revascularized by percutaneous coronary intervention. 2010 , 103, 512-21	191
	Value of a new multiparametric score for prediction of microvascular obstruction lesions in ST-segment elevation myocardial infarction revascularized by percutaneous coronary intervention.	
336	Value of a new multiparametric score for prediction of microvascular obstruction lesions in ST-segment elevation myocardial infarction revascularized by percutaneous coronary intervention. 2010, 103, 512-21 Clinical predictors of successful thrombectomy with the Export aspiration catheter in the acute	11
336 335	Value of a new multiparametric score for prediction of microvascular obstruction lesions in ST-segment elevation myocardial infarction revascularized by percutaneous coronary intervention. 2010, 103, 512-21 Clinical predictors of successful thrombectomy with the Export aspiration catheter in the acute phase of myocardial infarction. Data from the RICO survey working group. 2010, 103, 522-9	11
336 335 334	Value of a new multiparametric score for prediction of microvascular obstruction lesions in ST-segment elevation myocardial infarction revascularized by percutaneous coronary intervention. 2010, 103, 512-21 Clinical predictors of successful thrombectomy with the Export[] aspiration catheter in the acute phase of myocardial infarction. Data from the RICO survey working group. 2010, 103, 522-9 Successful reperfusion therapy: from epicardial to myocardial salvage. 2010, 63, 757-9 Myeloperoxidase may contribute to the no-reflow phenomenon in patients with acute myocardial	11 2
336 335 334 333	Value of a new multiparametric score for prediction of microvascular obstruction lesions in ST-segment elevation myocardial infarction revascularized by percutaneous coronary intervention. 2010, 103, 512-21 Clinical predictors of successful thrombectomy with the Export[] aspiration catheter in the acute phase of myocardial infarction. Data from the RICO survey working group. 2010, 103, 522-9 Successful reperfusion therapy: from epicardial to myocardial salvage. 2010, 63, 757-9 Myeloperoxidase may contribute to the no-reflow phenomenon in patients with acute myocardial infarction. 2010, 139, 187-92	11 2 1
336 335 334 333 332	Value of a new multiparametric score for prediction of microvascular obstruction lesions in ST-segment elevation myocardial infarction revascularized by percutaneous coronary intervention. 2010, 103, 512-21 Clinical predictors of successful thrombectomy with the Export[] aspiration catheter in the acute phase of myocardial infarction. Data from the RICO survey working group. 2010, 103, 522-9 Successful reperfusion therapy: from epicardial to myocardial salvage. 2010, 63, 757-9 Myeloperoxidase may contribute to the no-reflow phenomenon in patients with acute myocardial infarction. 2010, 139, 187-92 Unresolved issues in myocardial reperfusion injury. 2010, 19, 29-35 Angiographic assessment of microvascular perfusionmyocardial blush in clinical practice. 2010,	11 2 1 9 58

328	The assessment of microvascular flow and tissue perfusion using ultrasound imaging. 2010 , 224, 273-90	33
327	Microvascular occlusions and coronary microembolization. 2011 , 45, 258-60	
326	Coronary and microvascular thrombolysis with guided diagnostic ultrasound and microbubbles in acute ST segment elevation myocardial infarction. 2011 , 24, 1400-8	43
325	Value of speckle-tracking echocardiography for prediction of left ventricular remodeling in patients with ST-elevation myocardial infarction treated by primary percutaneous intervention. 2011 , 24, 1342-8	41
324	Myocardial contrast echocardiography in coronary artery disease. 2011 , 21, 1-11	1
323	Cardiac Magnetic Resonance Imaging in 1schemic Heart Disease. 2011 , 6, 453-73	
322	Protective effects of salvianolate on microvascular flow in a porcine model of myocardial ischaemia and reperfusion. 2011 , 104, 313-24	36
321	A multicenter, randomized, controlled study of mechanical left ventricular unloading with counterpulsation to reduce infarct size prepercutaneous coronary intervention for acute myocardial infarction: rationale and design of the Counterpulsation Reduces Infarct Size Acute	16
320	Effect of delayed vs immediate stent implantation on myocardial perfusion and cardiac function in patients with ST-segment elevation myocardial infarction undergoing primary percutaneous intervention with thrombus aspiration. 2011 , 27, 541-7	27
319	Usefulness of endothelin-1 assessment in acute myocardial infarction. 2011 , 64, 105-10	19
318	Usefulness of Endothelin-1 Assessment in Acute Myocardial Infarction. 2011 , 64, 105-110	3
317	Elevated plasma asymmetric dimethylarginine level in acute myocardial infarction patients as a predictor of poor prognosis and angiographic impaired reperfusion. 2011 , 219, 304-10	21
316	The association of serum uric acid levels on coronary flow in patients with STEMI undergoing primary PCI. 2011 , 219, 334-41	71
315	High admission levels of Eglutamyltransferase predict poor myocardial perfusion after primary percutaneous intervention. 2011 , 66, 1729-34	1
314	Microcirculation coronaire. 2011 , 6, 1-3	
313	Elevated serum uric acid predicts angiographic impaired reperfusion and 1-year mortality in ST-segment elevation myocardial infarction patients undergoing percutaneous coronary intervention. 2011 , 59, 931-7	24
312	Impact of obstructive sleep apnea on myocardial tissue perfusion in patients with ST-segment elevation myocardial infarction. 2011 , 75, 890-6	24
311	Distal protection during primary coronary intervention can preserve the index of microcirculatory resistance in patients with acute anterior ST-segment elevation myocardial infarction. 2011 , 75, 94-8	15

310	High QRS score on admission strongly predicts impaired myocardial reperfusion in patients with a first anterior acute myocardial infarction. 2011 , 75, 626-32	13
309	Long-term treatment with eicosapentaenoic acid ameliorates myocardial ischemia-reperfusion injury in pigs in vivoInvolvement of Rho-kinase pathway inhibition 2011 , 75, 1843-51	15
308	Ischemic Heart Disease. 2011 , 203-273	
307	Mechanical assistance by intra-aortic balloon pump counterpulsation during reperfusion increases coronary blood flow and mitigates the no-reflow phenomenon: an experimental study. 2011 , 35, 867-74	26
306	Integrilin in patients undergoing primary percutaneous coronary intervention for ST-elevation myocardial infarction. 2011 , 24, 351-6	9
305	Management of the no-reflow phenomenon. 2011 , 132, 72-85	21
304	New method of intracoronary adenosine injection to prevent microvascular reperfusion injury in patients with acute myocardial infarction undergoing percutaneous coronary intervention. 2011 , 107, 1131-5	39
303	[Usefulness of diastolic deceleration time assessed by transthoracic Doppler measurement in the detection of sustained microvascular obstruction in STEMI patients treated by primary PTCA]. 2011 , 60, 119-26	
302	A porcine model of ischemic heart failure produced by intracoronary injection of ethyl alcohol. 2011 , 26, 342-8	16
301	Comparison of the effects of nitroprusside versus nicorandil on the slow/no-reflow phenomenon during coronary interventions for acute myocardial infarction. 2011 , 26, 379-84	30
300	Relationship between retrograde coronary blood flow and the extent of no-reflow and infarct size in a porcine ischemia-reperfusion model. 2011 , 4, 99-105	1
299	The no-reflow phenomenon associated with percutaneous coronary intervention: its mechanisms and treatment. 2011 , 26, 2-11	8
298	A patient with significant slow-flow phenomenon during percutaneous coronary intervention for ST elevation myocardial infarction associated with scattered necrotic core by virtual histology intravascular ultrasound. 2011 , 26, 290-5	2
297	Non-invasive evaluation of myocardial reperfusion by transthoracic Doppler echocardiography and single-photon emission computed tomography in patients with anterior acute myocardial infarction. 2011 , 9, 16	3
296	The multidimensional physiological responses to postconditioning. 2011 , 14, 791-810	42
295	Ischemia/reperfusion injury and cardioprotective mechanisms: Role of mitochondria and reactive oxygen species. 2011 , 3, 186-200	218
294	High concentrations of asymmetric dimethylarginine are associated with ST-segment resolution failure after reperfusion for acute myocardial infarction. 2011 , 49, 903-7	5
293	Phosphorylation of endothelial NOS contributes to simvastatin protection against myocardial no-reflow and infarction in reperfused swine hearts: partially via the PKA signaling pathway. 2012 , 33, 879-87	23

292	Clinical features of early myocardial rupture of acute myocardial infarction. 2012, 63, 453-6	2
291	Why are We Interested in Viability?. 2012 , 155-171	
290	Impact of metabolic syndrome on various aspects of microcirculation and major adverse cardiac events in patients with ST-segment elevation myocardial infarction. 2012 , 76, 1972-9	16
289	Intracoronary hemodynamic effects of pressure-controlled intermittent coronary sinus occlusion (PICSO): results from the First-In-Man Prepare PICSO Study. 2012 , 25, 549-56	13
288	Five-year clinical outcomes after implantation of sirolimus-eluting stents in patients with and without diabetes mellitus. 2012 , 27, 189-95	7
287	Apport de lâIRM cardiaque dans le diagnostic des complications de lâInfarctus du myocarde. 2012 , 93, 611-619	
286	The role of cardiovascular imaging to understand the different patterns of post-ischemic remodeling. 2012 , 22, 107-117	
285	A proinflammatory monocyte response is associated with myocardial injury and impaired functional outcome in patients with ST-segment elevation myocardial infarction: monocytes and myocardial infarction. 2012 , 163, 57-65.e2	83
284	Relationship between microcirculatory dysfunction and resolution of ST-segment elevation in the early phase after primary angioplasty in patients with ST-segment elevation myocardial infarction. 2012 , 159, 144-9	5
283	Remodeling after acute myocardial infarction: mapping ventricular dilatation using three dimensional CMR image registration. 2012 , 14, 41	20
282	CMR of microvascular obstruction and hemorrhage in myocardial infarction. 2012 , 14, 68	112
281	Cardiac MRI in the diagnosis of complications of myocardial infarction. 2012 , 93, 578-85	6
280	Differences in clinical and angiographic outcomes with different drug-eluting stents in Japanese patients with and without diabetes mellitus. 2012 , 60, 361-6	11
279	Imaging the myocardial microcirculation post-myocardial infarction. 2012 , 9, 282-92	13
278	Index of microcirculatory resistance as predictor for microvascular functional recovery in patients with anterior myocardial infarction. 2012 , 27, 1044-50	17
277	Elevated admission microalbuminuria predicts poor myocardial blood flow and 6-month mortality in ST-segment elevation myocardial infarction patients undergoing primary percutaneous coronary intervention. 2012 , 35, 219-24	9
276	Myocardial contrast echocardiography after myocardial infarction. 2012 , 14, 350-8	5
275	Why Quantify Myocardial Perfusion?. 2012 , 5, 133-143	4

274	Implications of ST-segment elevation in leads V5 and V6 in patients with reperfused inferior wall acute myocardial infarction. 2012 , 109, 314-9		4
273	Positron emission tomography in acute coronary syndromes. 2012 , 5, 11-21		2
272	Contrast media injection into right coronary artery caused thrombus extraction to aorta that may have resulted in left main trunk thrombosis. 2012 , 27, 37-42		
271	Safety and feasibility of high-dose administration of nicorandil before reperfusion therapy in acute myocardial infarction. 2013 , 28, 352-61		9
270	Beneficial effects of intracoronary nicorandil on microvascular dysfunction after primary percutaneous coronary intervention: demonstration of its superiority to nitroglycerin in a cross-over study. <i>Cardiovascular Drugs and Therapy</i> , 2013 , 27, 279-87	3.9	32
269	Acute myocardial infarction: early CT aspects of myocardial microcirculation obstruction after percutaneous coronary intervention. 2013 , 23, 2405-12		8
268	Myocardial reperfusion injury: looking beyond primary PCI. 2013 , 34, 1714-22		252
267	Prognostic value of microvascular resistance index immediately after primary percutaneous coronary intervention on left ventricular remodeling in patients with reperfused anterior acute ST-segment elevation myocardial infarction. 2013 , 6, 1046-54		25
266	Novel MRI and CT Approaches for the Characterization of Myocardial Infarct. 2013, 1, 233-245		
265	Magnetic resonance imaging-defined areas of microvascular obstruction after acute myocardial infarction represent microvascular destruction and haemorrhage. 2013 , 34, 2346-53		141
264	The effect of prethrombolytic cyclosporine-A injection on clinical outcome of acute anterior ST-elevation myocardial infarction. 2013 , 31, e34-9		45
263	Chronic pretreatment of metformin is associated with the reduction of the no-reflow phenomenon in patients with diabetes mellitus after primary angioplasty for acute myocardial infarction. 2013 , 31, 60-4		16
262	Impact of attenuated plaque as detected by intravascular ultrasound on the occurrence of microvascular obstruction after percutaneous coronary intervention in patients with ST-segment elevation myocardial infarction. 2013 , 6, 847-53		31
261	Impact of thrombus aspiration during primary percutaneous coronary intervention in cardiogenic shock complicating ST-segment elevation myocardial infarction. 2013 , 14, 307-10		5
2 60	The role of cardiac magnetic resonance imaging (MRI) in acute myocardial infarction (AMI). 2013 , 22, 243-55		26
259	Longitudinal extent of lipid pool assessed by optical coherence tomography predicts microvascular no-reflow after primary percutaneous coronary intervention for ST-segment elevation myocardial infarction. 2013 , 62, 71-6		21
258	Clinical, angiographic, and intravascular ultrasound predictors of early stent thrombosis in patients with acute myocardial infarction. 2013 , 168, 1674-5		6
257	Estimation of infarct size using transthoracic Doppler echocardiographic measurement of coronary flow reserve in infarct related and reference coronary artery. 2013 , 168, 169-75		6

256	Impaired coronary flow reserve after a recent myocardial infarction: correlation with infarct size and extent of microvascular obstruction. 2013 , 167, 351-6	17
255	Degenerated Saphenous Vein Graft Intervention: Should We Target the Native Vessel instead?. 2013 , 2, 323-337	2
254	Use of thrombectomy devices in primary percutaneous coronary intervention: a systematic review and meta-analysis. 2013 , 163, 229-241	46
253	Development and validation of a clinical risk score predicting the no-reflow phenomenon in patients treated with primary percutaneous coronary intervention for ST-segment elevation myocardial infarction. 2013 , 124, 153-60	15
252	Effect of new method of intracoronary adenosine injection during primary percutaneous coronary intervention on microvascular reperfusion injury - clinical outcome and 1-year follow-up. 2013 , 124, 199-206	4
251	Thrombectomy Devices in Coronary Intervention. 2013 , 2, 347-359	
250	Innate immune response after acute myocardial infarction and pharmacomodulatory action of tacrolimus in reducing infarct size and preserving myocardial integrity. 2013 , 20, 82	17
249	Detection of acute reperfusion myocardial hemorrhage with cardiac MR imaging: T2 versus T2. 2013 , 269, 387-95	62
248	Glucagon-like peptide-1 preserves coronary microvascular endothelial function after cardiac arrest and resuscitation: potential antioxidant effects. 2013 , 304, H538-46	17
247	Methods to investigate coronary microvascular function in clinical practice. 2013 , 14, 1-18	45
247 246	Methods to investigate coronary microvascular function in clinical practice. 2013 , 14, 1-18 Myocardial ischemia-reperfusion injury: a neglected therapeutic target. 2013 , 123, 92-100	1250
246	Myocardial ischemia-reperfusion injury: a neglected therapeutic target. 2013 , 123, 92-100 Effect of statin pretreatment on myocardial perfusion in patients undergoing primary	1250
246 245	Myocardial ischemia-reperfusion injury: a neglected therapeutic target. 2013 , 123, 92-100 Effect of statin pretreatment on myocardial perfusion in patients undergoing primary percutaneous coronary intervention: a systematic review and meta-analysis. 2013 , 36, E17-24	1250
246 245 244	Myocardial ischemia-reperfusion injury: a neglected therapeutic target. 2013 , 123, 92-100 Effect of statin pretreatment on myocardial perfusion in patients undergoing primary percutaneous coronary intervention: a systematic review and meta-analysis. 2013 , 36, E17-24 Rationale and design of the MGuard for acute ST elevation reperfusion MASTER trial. 2013 , 82, 184-90	1250 6 7
246245244243	Myocardial ischemia-reperfusion injury: a neglected therapeutic target. 2013, 123, 92-100 Effect of statin pretreatment on myocardial perfusion in patients undergoing primary percutaneous coronary intervention: a systematic review and meta-analysis. 2013, 36, E17-24 Rationale and design of the MGuard for acute ST elevation reperfusion MASTER trial. 2013, 82, 184-90 Primary percutaneous coronary intervention: devices to prevent no-reflow phenomenon. 2013, 5, 289-300 Application of adenosine stress echocardiography in the prognosis of acute myocardial infarction	1250 6 7 2
246245244243242	Myocardial ischemia-reperfusion injury: a neglected therapeutic target. 2013, 123, 92-100 Effect of statin pretreatment on myocardial perfusion in patients undergoing primary percutaneous coronary intervention: a systematic review and meta-analysis. 2013, 36, E17-24 Rationale and design of the MGuard for acute ST elevation reperfusion MASTER trial. 2013, 82, 184-90 Primary percutaneous coronary intervention: devices to prevent no-reflow phenomenon. 2013, 5, 289-300 Application of adenosine stress echocardiography in the prognosis of acute myocardial infarction following percutaneous coronary interventional therapy. 2013, 6, 727-730 Thrombus aspiration therapy and coronary thrombus components in patients with acute	1250 6 7 2

238	Inhibition of dipeptidyl peptidase-IV enzyme activity protects against myocardial ischemia-reperfusion injury in rats. 2014 , 12, 357	28
237	Uric acid as a predictor of in-hospital mortality in acute myocardial infarction: a meta-analysis. 2014 , 70, 1597-601	15
236	Complement factor concentrations in patients with acute myocardial infarction: time course and ability to predict left ventricular dysfunction. 2014 , 8, 247-58	1
235	Clinical importance of hyperemic coronary blood flow (thrombolysis in myocardial infarction-intravenous flow) after primary percutaneous coronary intervention. 2014 , 25, 665-70	
234	Does Serum Bilirubin level on admission predict TIMI flow grade and in-hospital MACE in patients with STEMI undergoing primary PCI. 2014 , 65, 198-204	44
233	The central role of conventional 12-lead ECG for the assessment of microvascular obstruction after percutaneous myocardial revascularization. 2014 , 47, 45-51	12
232	Impact of nonculprit vessel myocardial perfusion on outcomes of patients undergoing percutaneous coronary intervention for acute coronary syndromes: analysis from the ACUITY trial (Acute Catheterization and Urgent Intervention Triage Strategy). 2014 , 7, 266-75	2
231	The "no reflow" phenomenon following acute myocardial infarction: mechanisms and treatment options. 2014 , 64, 77-85	44
230	Impact of runoff grade after endovascular therapy for femoropopliteal lesions. 2014 , 59, 720-7	19
229	CMR assessment of microvascular obstruction in STEMI: ready for prime time?. 2014 , 64, 1227-30	5
228	Microvascular obstruction in patients with non-ST-elevation myocardial infarction: a contrast-enhanced cardiac magnetic resonance study. 2014 , 30, 1087-95	12
227	Evidence of myocardial scarring and microvascular obstruction on cardiac magnetic resonance imaging in a series of patients presenting with myocardial infarction without obstructed coronary arteries. 2014 , 30, 1097-103	4
226	Prognostic value of microvascular obstruction and infarct size, as measured by CMR in STEMI patients. 2014 , 7, 930-9	195
225	Influence of microvascular obstruction on regional myocardial deformation in the acute phase of myocardial infarction: a speckle-tracking echocardiography study. 2014 , 27, 93-100	16
224	Relation of ST-segment elevation before and after percutaneous transluminal coronary angioplasty to left ventricular area at risk, myocardial infarct size, and systolic function. 2014 , 113, 593-600	14
223	No-reflow phenomenon: pathophysiology, diagnosis, prevention, and treatment. A review of the current literature and future perspectives. 2014 , 65, 180-9	51
222	Association of lymphocyte-to-monocyte ratio with the no-reflow phenomenon in patients who underwent a primary percutaneous coronary intervention for ST-elevation myocardial infarction. 2015 , 26, 706-12	38
221	QRS duration: a novel marker of microvascular reperfusion as assessed by myocardial blush grade in ST elevation myocardial infarction patients undergoing a primary percutaneous intervention. 2015 , 26, 583-6	6

(2015-2015)

220	Meta-Analysis of Randomized Controlled Trials. 2015 , 2015, 382086	21
219	Prognostic Value of Speckle Tracking Echocardiography in Patients with ST-Elevation Myocardial Infarction Treated with Late Percutaneous Intervention. 2015 , 32, 1384-91	22
218	Effect of infarct severity on regional and global left ventricular remodeling in patients with successfully reperfused ST segment elevation myocardial infarction. 2015 , 274, 93-102	20
217	Late gadolinium enhancement imaging in assessment of myocardial viability: techniques and clinical applications. 2015 , 53, 397-411	22
216	Treatment of microvascular micro-embolization using microbubbles and long-tone-burst ultrasound: an in vivo study. 2015 , 41, 456-64	24
215	Cellular postconditioning: allogeneic cardiosphere-derived cells reduce infarct size and attenuate microvascular obstruction when administered after reperfusion in pigs with acute myocardial infarction. 2015 , 8, 322-32	65
214	Myocardial infarction. 2015 , 271-325	
213	Predictive Factors of Pericardial Effusion After a First Acute Myocardial Infarction and Successful Reperfusion. 2015 , 116, 497-503	3
212	Novel therapeutics in myocardial infarction: targeting microvascular dysfunction and reperfusion injury. 2015 , 36, 605-16	40
211	Microvascular obstruction on delayed enhancement cardiac magnetic resonance imaging after acute myocardial infarction, compared with myocardial (201)Tl and (123)I-BMIPP dual SPECT findings. 2015 , 84, 1516-1524	2
210	Thrombus aspiration in patients with ST elevation myocardial infarction: meta-analysis of 16 randomized trials. 2015 , 15, 175-87	3
209	Association between hyperglycemia at admission and microvascular obstruction in patients with ST-segment elevation myocardial infarction. 2015 , 65, 272-7	14
208	Novel insights into an "old" phenomenon: the no reflow. 2015 , 187, 273-80	76
207	Ischemic preconditioning-an unfulfilled promise. 2015 , 16, 101-8	7
206	Soluble interleukin 6 receptor levels are associated with reduced myocardial reperfusion after percutaneous coronary intervention for acute myocardial infarction. 2015 , 73, 207-12	11
205	Use and impact of thrombectomy in primary percutaneous coronary intervention for acute myocardial infarction with persistent ST-segment elevation: results of the prospective ALKK PCI-registry. 2015 , 104, 803-11	7
204	Comparison of the improvement in myocardial perfusion and function in cases of rapid and slow electrocardiographic stage dynamics between patients with TIMI-3 flow after primary angioplasty for acute myocardial infarction. 2015 , 51, 217-21	1
203	The no-reflow phenomenon: State of the art. 2015 , 108, 661-74	90

202	[No reflow: What are the predictors?]. 2015 , 64, 472-80	1
201	Association of nonculprit plaque characteristics with transient slow flow phenomenon during percutaneous coronary intervention. 2015 , 181, 108-13	6
200	Resting qualitative and quantitative myocardial contrast echocardiography to predict cardiac events in patients with acute myocardial infarction and percutaneous revascularization. 2015 , 30, 45-55	8
199	Treating thrombus in the coronary arteries. 2015 , 40, 60-5	2
198	Intramyocardial haemorrhage after acute myocardial infarction. 2015, 12, 156-67	88
197	Serial assessments of microvascular obstruction by contrast-enhanced magnetic resonance predict contractile recovery and clinical outcome after reperfused acute myocardial infarction. 2015 , 62, 345-57	3
196	No-Reflow Phoenomenon by Intracoronary Thrombus in Acute Myocardial Infarction. 2016 , 52, 38-44	7
195	The association of serum procalcitonin level with the no-reflow phenomenon after a primary percutaneous coronary intervention in patients with ST-elevation myocardial infarction. 2016 , 27, 116-21	6
194	Concordance of angiographic and electrocardiographic indexes of microvascular obstruction: myocardial haemorrhage role. 2016 , 17, 382-91	2
193	Prognostic utility of myocardial blush grade after PCI in patients with NSTE-ACS: Analysis from the ACUITY trial. 2016 , 88, 215-24	5
192	Coronary Artery Dissections, Perforations, and the No-Reflow Phenomenon. 2016 , 248-266	
191	Fragmented QRS complexes have predictive value of imperfect ST-segment resolution in patients with STEMI after primary percutaneous coronary intervention. 2016 , 34, 398-402	8
190	Efficacy and Safety of Local Intracoronary Drug Delivery in Treatment of No-Reflow Phenomenon: A Pilot Study. 2016 , 29, 496-504	7
189	Growth differentiation factor 15 may protect the myocardium from no-reflow by inhibiting the inflammatory-like response that predominantly involves neutrophil infiltration. 2016 , 13, 623-32	9
188	Safety and feasibility of local myocardial hypothermia. 2016 , 87, 877-83	9
187	Effect of Thrombus Composition and Viscosity on Sonoreperfusion Efficacy in a Model of Micro-Vascular Obstruction. 2016 , 42, 2220-31	7
186	Comparison between visual grading and planimetric quantification of microvascular obstruction extent assessment in reperfused acute myocardial infarction. 2016 , 26, 2166-75	4
185	Evaluating the Use of Spatial Covariance Structures in the Analysis of Cardiovascular Imaging Data. 2016 , 45, 3149-3170	1

184	Amelioration of persistent left ventricular function impairment through increased plasma ascorbate levels following myocardial infarction. 2016 , 21, 75-83	14
183	The challenges and impact of microvascular injury in ST-elevation myocardial infarction. 2016 , 14, 431-43	21
182	Facing Time in Ischemic Stroke: An Alternative Hypothesis for Collateral Failure. 2016 , 26, 141-51	29
181	Thrombus aspiration in acute myocardial infarction. 2016 , 13, 418-28	27
180	Morphological predictors for no reflow phenomenon after primary percutaneous coronary intervention in patients with ST-segment elevation myocardial infarction caused by plaque rupture. 2017 , 18, 103-110	31
179	ST-segment resolution with bivalirudin versus heparin and routine glycoprotein IIb/IIIa inhibitors started in the ambulance in ST-segment elevation myocardial infarction patients transported for primary percutaneous coronary intervention: The EUROMAX ST-segment resolution substudy. 2017	4
178	Coronary flow improvement following unsuccessful primary percutaneous coronary intervention in ST-elevation myocardial infarction with diffuse ectatic coronary artery. 2017 , 6, 623-631	7
177	Impact of thermodilution-derived coronary blood flow patterns after percutaneous coronary intervention on mid-term left ventricular remodeling in patients with ST elevation myocardial infarction. 2017 , 32, 1-7	1
176	Plasma big endothelin-1 levels at admission and future cardiovascular outcomes: A cohort study in patients with stable coronary artery disease. 2017 , 230, 76-79	18
175	Predictors of no- reflow during primary angioplasty for acute myocardial infarction, from Medical College Hospital, Trivandrum. 2017 , 69 Suppl 1, S34-S45	14
174	Lymphocyte Communication in Myocardial Ischemia/Reperfusion Injury. 2017, 26, 660-675	38
173	Current perspectives in coronary microvascular dysfunction. 2017 , 24, e12340	24
172	Clinical role of post-angioplasty hyperemic microvascular resistances in chronic ischemic left ventricular dysfunction. 2017 , 18, 332-340	3
171	Usefulness of Mean Platelet Volume-to-Lymphocyte Ratio for Predicting Angiographic No-Reflow and Short-Term Prognosis After Primary Percutaneous Coronary Intervention in Patients With ST-Segment Elevation Myocardial Infarction. 2017 , 120, 534-541	25
170	Salvianolate reduces murine myocardial ischemia and reperfusion injury via ERK1/2 signaling pathways in vivo. 2017 , 23, 40-47	15
169	Microvascular Obstruction in Acute Myocardial Infarction. 2017 , 3, 197-202	
168	The Role of Nitric Oxide during Sonoreperfusion of Microvascular Obstruction. 2017 , 7, 3527-3538	15
167	Value of 3-Dimensional Speckle Tracking Echocardiography in the Prediction of Microvascular Obstruction and Left Ventricular Remodeling in Patients With ST-Elevation Myocardial Infarction. 2017 , 81, 353-360	7

166	Prevalence and Predictive Value of Microvascular Flow Abnormalities after Successful Contemporary Percutaneous Coronary Intervention in Acute ST-Segment Elevation Myocardial Infarction. 2018 , 31, 674-682		14
165	Which admission electrocardiographic parameter is more powerful predictor of no-reflow in patients with acute anterior myocardial infarction who underwent primary percutaneous intervention?. 2018 , 51, 203-209		5
164	Cardiosphere-Derived Cells and Ischemic Heart Failure. 2018 , 26, 8-21		29
163	MiR-124 aggravates failing hearts by suppressing CD151-facilitated angiogenesis in heart. 2018 , 9, 143	382-143	8 96 6
162	Molecular Imaging of VWF (von Willebrand Factor) and Platelet Adhesion in Postischemic Impaired Microvascular Reflow. 2018 , 11, e007913		10
161	Prognostic significance of fragmented QRS in patients with ST-elevation myocardial infarction undergoing revascularization. 2018 , 70 Suppl 3, S126-S132		5
160	Combined Therapy with SS31 and Mitochondria Mitigates Myocardial Ischemia-Reperfusion Injury in Rats. 2018 , 19,		32
159	Strain analysis is superior to wall thickening in discriminating between infarcted myocardium with and without microvascular obstruction. 2018 , 28, 5171-5181		12
158	Epidemiology of Coronary Microvascular Obstruction. 2018, 53-68		
157	Role of Ischemiaâ R eperfusion Injury in Coronary MVO. 2018 , 97-107		
157 156	Role of IschemiaâReperfusion Injury in Coronary MVO. 2018, 97-107 Imaging Modalities for the Assessment of CMVO. 2018, 155-166		
156	Imaging Modalities for the Assessment of CMVO. 2018 , 155-166 Long-Term Management After Coronary Microvascular Obstruction Complicating Reperfusion in		10
156 155	Imaging Modalities for the Assessment of CMVO. 2018, 155-166 Long-Term Management After Coronary Microvascular Obstruction Complicating Reperfusion in ST-Elevation Myocardial Infarction. 2018, 341-357 Microvascular obstruction in non-infarct related coronary arteries is an independent predictor of major adverse cardiovascular events in patients with ST segment-elevation myocardial infarction.		10
156 155 154	Imaging Modalities for the Assessment of CMVO. 2018, 155-166 Long-Term Management After Coronary Microvascular Obstruction Complicating Reperfusion in ST-Elevation Myocardial Infarction. 2018, 341-357 Microvascular obstruction in non-infarct related coronary arteries is an independent predictor of major adverse cardiovascular events in patients with ST segment-elevation myocardial infarction. 2018, 273, 22-28 ROUTINELY AVAILABLE BIOMARKERS AS LONG-TERM PREDICTORS OF DEVELOPING SYSTOLIC DYSFUNCTION IN COMPLETELY REVASCULARIZED PATIENTS WITH ACUTE ST ELEVATION		
156 155 154	Imaging Modalities for the Assessment of CMVO. 2018, 155-166 Long-Term Management After Coronary Microvascular Obstruction Complicating Reperfusion in ST-Elevation Myocardial Infarction. 2018, 341-357 Microvascular obstruction in non-infarct related coronary arteries is an independent predictor of major adverse cardiovascular events in patients with ST segment-elevation myocardial infarction. 2018, 273, 22-28 ROUTINELY AVAILABLE BIOMARKERS AS LONG-TERM PREDICTORS OF DEVELOPING SYSTOLIC DYSFUNCTION IN COMPLETELY REVASCULARIZED PATIENTS WITH ACUTE ST ELEVATION MYOCARDIAL INFARCTION. 2019, 58, 95-102 Relationship between the ST-Segment Resolution and Microvascular Dysfunction in Patients Who	2.4	2
156 155 154 153	Imaging Modalities for the Assessment of CMVO. 2018, 155-166 Long-Term Management After Coronary Microvascular Obstruction Complicating Reperfusion in ST-Elevation Myocardial Infarction. 2018, 341-357 Microvascular obstruction in non-infarct related coronary arteries is an independent predictor of major adverse cardiovascular events in patients with ST segment-elevation myocardial infarction. 2018, 273, 22-28 ROUTINELY AVAILABLE BIOMARKERS AS LONG-TERM PREDICTORS OF DEVELOPING SYSTOLIC DYSFUNCTION IN COMPLETELY REVASCULARIZED PATIENTS WITH ACUTE ST ELEVATION MYOCARDIAL INFARCTION. 2019, 58, 95-102 Relationship between the ST-Segment Resolution and Microvascular Dysfunction in Patients Who Underwent Primary Percutaneous Coronary Intervention. 2019, 2019, 8695065 Impact of intracoronary contrast injection pressure on reperfusion during primary percutaneous coronary intervention in acute ST-segment elevation myocardial infarction: A prospective	2.4	2

148	Imaging the injured beating heart intravitally and the vasculoprotection afforded by haematopoietic stem cells. 2019 , 115, 1918-1932	11
147	Multilayer longitudinal strain can help predict the development of no-reflow in patients with acute coronary syndrome without ST elevation. 2019 , 35, 1811-1821	2
146	The cyclooxygenase-1/mPGES-1/endothelial prostaglandin EP4 receptor pathway constrains myocardial ischemia-reperfusion injury. 2019 , 10, 1888	30
145	Comparison of 0.9-mm and 1.4-mm catheters in excimer laser coronary angioplasty for acute myocardial infarction. 2019 , 34, 1747-1754	5
144	Primary percutaneous coronary intervention in ST-elevation myocardial infarction with an ectatic infarct-related artery. 2019 , 30, 277-284	
143	Efficacy and safety of thrombus aspiration in ST-segment elevation myocardial infarction: an updated systematic review and meta-analysis of randomised clinical trials. 2019 , 8, 24-38	7
142	The Effect of Stent Artefact on Quantification of Plaque Features Using Optical Coherence Tomography (OCT): A Feasibility and Clinical Utility Study. 2020 , 29, 874-882	2
141	Incidence, predictors and outcomes of stress hyperglycemia in patients with ST elevation myocardial infarction undergoing primary percutaneous coronary intervention. 2020 , 17, 1479164119883983	12
140	Non-infarct related artery microvascular obstruction is associated with worse persistent diastolic dysfunction in patients with revascularized ST elevation myocardial infarction. 2020 , 300, 27-33	4
139	The CHADS-VASc score is a predictor of no-reflow in patients with non-ST-segment elevation myocardial infarction. 2020 , 31, 7-12	12
138	No-Reflow after PPCI-A Predictor of Short-Term Outcomes in STEMI Patients. 2020 , 9,	7
137	All-cause mortality and the risk of stroke with selective aspiration thrombectomy in patients with ST-elevation myocardial infarction undergoing primary percutaneous coronary intervention: A nationwide retrospective cohort study. 2020 , 99, e19590	1
136	Assessment of Growth Differentiation Factor 15 Levels on Coronary Flow in Patients with STEMI Undergoing Primary PCI. 2020 , 8,	0
135	Sonoreperfusion therapy for microvascular obstruction: A step toward clinical translation. 2020 , 46, 712-720	2
134	Incidence, Predictors, and Prognosis of Coronary Slow-Flow and No-Reflow Phenomenon in Patients with Chronic Total Occlusion Who Underwent Percutaneous Coronary Intervention. 2020 , 16, 95-101	4
133	Predictive values of D-dimer for the long-term prognosis of acute ST-segment elevation infarction: A retrospective study in southwestern China. 2020 , 99, e19724	2
132	A Scoring System to Predict No-Reflow Phenomenon in Elective Percutaneous Coronary Intervention: The RECOVER Score. 2021 , 46, 100676	
131	Prognostic value of initial QRS analysis in anterior STEMI: Correlation with left ventricular systolic dysfunction, serum biomarkers, and cardiac outcomes. 2021 , 26, e12791	2

130	Low Ankle-Brachial Index is Associated with Microvascular Coronary Obstruction After Primary PCI. 2021 , 17, 23-32		
129	Systematic Understanding of Pathophysiological Mechanisms of Oxidative Stress-Related Conditions-Diabetes Mellitus, Cardiovascular Diseases, and Ischemia-Reperfusion Injury. <i>Frontiers in Cardiovascular Medicine</i> , 2021 , 8, 649785	5.4	8
128	The outcomes in STEMI patients with high thrombus burden treated by deferred versus immediate stent implantation in primary percutaneous coronary intervention: a prospective cohort study. 2021 , 9, 573		1
127	Enhancement patterns detected by multidetector computed tomography are associated with the long-term prognosis of patients with acute myocardial infarction. 2021 , 36, 1784-1793		
126	In Search of the Perfect Coronary Perfusion. 2021 , 116, 865-866		
125	Predictors of no-reflow phenomenon following percutaneous coronary intervention for ST-segment elevation myocardial infarction. 2021 , 70, 136-142		2
124	Reactive Hyperglycaemia Connected with Low Glycated Haemoglobin âlRisk Factor for Cardiovascular Adverse Events. 2021 , 2, 85-96		
123	Accuracy of Post-thrombolysis ST-segment Reduction as an Adequate Reperfusion Predictor in the Pharmaco-Invasive Approach. 2021 , 117, 15-25		О
122	Metformin attenuates ischemia/reperfusion-induced apoptosis of cardiac cells by downregulation of p53/microRNA-34a via activation of SIRT1. 2021 , 99, 875-884		O
121	Controlling Reperfusion Injury With Controlled Reperfusion: Historical Perspectives and New Paradigms. 2021 , 26, 504-523		4
120	How to Mediate Cardioprotection in Ischemic heartsâ R ole of Adenosine. 2003 , 293-316		1
119	Does Reperfusion Cause Any Injury to the Myocardium?. 2003 , 145-157		1
118	Insights into the Pathophysiology of Acute Coronary Syndromes Using the TIMI Flow Grade and TIMI Frame Counting Methods. <i>Contemporary Cardiology</i> , 1999 , 87-107	0.1	1
117	Cardiovascular Disease and Magnesium. 2007 , 227-238		2
116	Intravenous myocardial contrast echocardiography for myocardial perfusion. <i>Developments in Cardiovascular Medicine</i> , 1996 , 461-472		1
115	Reduced microvascular and myocardial damage in patients with acute myocardial infarction and preinfarction angina. 2002 , 144, 796-803		32
114	Acute Coronary Syndromes. 2009 , 535-596		9
113	Medical and surgical therapy for cardiac remodeling. 1999 , 14, 196-205		6

112	Magnetic resonance imaging in acute myocardial infarction. 1999 , 14, 480-4		5
111	Unmet goals in the treatment of Acute Myocardial Infarction: Review. 2017, 6,		5
110	No-Reflow Phenomenon in Central Retinal Artery Occlusion: Incidence, Risk Factors, and Clinical Implications. <i>PLoS ONE</i> , 2015 , 10, e0142852	3.7	10
109	The value of real-time myocardial contrast echocardiography for detecting coronary microcirculation function in coronary artery disease patients. 2018 , 19, 27-33		2
108	Relationship between the mismatch of 123I-BMIPP and 201Tl myocardial single-photon emission computed tomography and autonomic nervous system activity in patients with acute myocardial infarction. 2006 , 47, 193-207		1
107	Aspiration thrombectomy of a massive thrombotic embolus in acute myocardial infarction caused by coronary embolism. 2007 , 48, 387-92		24
106	Effect of adjunctive tirofiban therapy on angiographic and clinical outcomes in patients with ST-segment elevated acute myocardial infarction undergoing primary stenting. 2004 , 45, 31-41		6
105	Pharmacological prevention of peri-, and post-procedural myocardial injury in percutaneous coronary intervention. 2008 , 4, 223-30		5
104	Anti-ischemic and antiarrhythmic activity of dibornol in conditions of multiple transitory myocardial ischemia in rats. 2020 , 35, 151-158		O
103	Assessment of No-Reflow Phenomenon by Myocardial Blush Grade and Pulsed Wave Tissue Doppler Imaging in Patients with Acute Coronary Syndrome. 2014 , 24, 52-56		1
102	The myocardial microcirculation: A key target for salvaging ischemic myocardium?. 2013, 03, 8-16		1
101	Bivalirudin infusion to reduce ventricular infarction: the open-label, randomised Bivalirudin Infusion for Ventricular InfArction Limitation (BIVAL) study. <i>EuroIntervention</i> , 2017 , 13, e540-e548	3.1	5
100	Systematic primary aspiration in acute myocardial percutaneous intervention: a multicentre randomised controlled trial of the export aspiration catheter. <i>EuroIntervention</i> , 2008 , 4, 222-8	3.1	55
99	Coronary thrombectomy, technical comparison of two systems on a laboratory bench: the impact of bends, angles and thrombus age. <i>EuroIntervention</i> , 2011 , 6, 729-34	3.1	4
98	Pre-infarction angina predicts thrombus burden in patients admitted for ST-segment elevation myocardial infarction. <i>EuroIntervention</i> , 2012 , 7, 1396-405	3.1	17
97	A new electrocardiographic marker of myocardial reperfusion in patients with acute ST-segment elevation myocardial infarction treated with primary percutaneous intervention: the value of QRS duration. <i>EuroIntervention</i> , 2012 , 7, 1406-12	3.1	4
96	Deferred stent implantation in patients with ST-segment elevation myocardial infarction: a pilot study. <i>EuroIntervention</i> , 2013 , 8, 1126-33	3.1	29
95	Reperfusion injury components and manifestations determined by cardiovascular MR and MDCT imaging. 2010 , 2, 1-14		6

94	Evaluation of the Clinical and Procedural Predictive Factors of no-Reflow Phenomenon Following Primary Percutaneous Coronary Intervention. <i>Research in Cardiovascular Medicine</i> , 2015 , 4, e25414	0.4	17
93	Predictors of in-hospital mortality in patients with ST-segment elevation myocardial infarction undergoing pharmacoinvasive treatment. 2013 , 68, 1516-20		16
92	Imaging Stem Cell-Based Myocardial Vasculoprotection. 2021 , 1-26		
91	Mechanism and potential treatment of the "no reflow" phenomenon after acute myocardial infarction: role of pericytes and GPR39. 2021 , 321, H1030-H1041		1
90	Role of NLRP3 Inflammasome in Myocardial Ischemia-Reperfusion Injury and Ventricular Remodeling 2022 , 28, e934255		1
89	??????????????? : ????????(?63? ?????????). Journal of JCS Cardiologists, 2000 , 8, 81-87	0.1	
88	Carbon Monoxide and Cardiovascular Inflammation. 2001 , 165-180		
87	Enhancement of Coronary Blood Flow as Myocardial Salvage Therapy. 2002 , 178-190		
86	Stress Echocardiography Versus Cardiac Magnetic Resonance Imaging. 2003, 403-418		
85	Distal Protection Devices. 2003 , 139-144		
84	Modulation of cardiac remodeling by adenosine: In vitro and in vivo effects. 2003, 17-26		
83	How Does Adenosine Mediate Cardioprotection?. Developments in Cardiovascular Medicine, 2003, 439-4	155	
82	Assessing Reperfusion and Prognostic Infarct Sizing with Biochemical Markers. 2003, 59-86		
81	Quantitative Analysis Using Overlap Images with Technetium-99m Pyrophosphate Thallium-201 Dual-isotope SPECT to Evaluate Microvascular Reperfusion Injury Following Successful Coronary Intervention in Patients with Acute Myocardial Infarction. 2004 , 16, 117-126		
80	Pathophysiological basics of acute myocardial infarction The epicardial vessel The microcirculation The myocardium. 2004 , 12-23		
79	Pathophysiological basics of acute myocardial infarction. 2004 , 3-14		
78	Complications and how to deal with them. 2004 , 57-74		2
77	Right heart catheterization and hemodynamic profiles. 2004 , 178-231		

76	Plaque Rupture: Pathoanatomical and Biomechanical Considerations. 2004, 73-90			
75	Differences Between Unstable Angina and Acute Myocardial Infarction: Pathophysiological and Clinical Spectrum. 2004 , 143-170			
74	Cardiac Rupture: Pathobiology, Diagnosis, Medical Management, and Surgical Intervention. 2004 , 597	-628		
73	Tissue Inflammation Impairs Tissue-Level Perfusion and Promotes Left Ventricular Remodeling in Patients With Acute Myocardial Infarction. <i>Journal of Echocardiography</i> , 2005 , 3, 83-90	1.6		
72	Vascular smooth muscle cell microparticles and coronary no-reflow: another piece of the puzzle?. <i>Thrombosis and Haemostasis</i> , 2005 , 94, 693-4	7		
71	Quantitative Myocardial Contrast Echocardiography in Coronary Artery Disease. <i>Journal of Echocardiography</i> , 2006 , 4, 99-109	1.6	1	
70	Assessment of Myocardial Perfusion With Intravenous Myocardial Contrast Echocardiography Current State and Clinical Applications. <i>Journal of Echocardiography</i> , 2006 , 4, 91-98	1.6		
69	Abrupt vessel closure and no-reflow. 2006 , 85-101			
68	Correlation between Thrombolysis in Myocardial Infarction, the Myocardial Perfusion Grade and the Myocardial Viability Indices after Primary Percutaneous Coronary Intervention in ST Segment Elevation Myocardial Infarction. <i>Korean Circulation Journal</i> , 2007 , 37, 581	2.2	1	
67	Successful Myocardial Reperfusion. Fundamental and Clinical Cardiology, 2008, 29-41			
66	Rescue Coronary Intervention for Failed Thrombolysis. 2009 , 101-113			
65	Myocardial Contrast Echocardiography in the Emergency Department. 2009, 415-425			
64	Pathology of and therapeutic strategy for microvascular dysfunction in patients with acute myocardial infarction. <i>Okayama Igakkai Zasshi</i> , 2009 , 121, 91-98	0		
63	Pathophysiology Basics of Acute Myocardial Infarction. 2009 , 1-14			
62	Major Subgroups at Presentation. 2009 , 15-44			
61	Major Subgroups at Presentation. 2009 , 15-44			
60	Myocardial Viability. 2010 , 267-283			
59	Novos parfinetros ecocardiogrficos para anlise da fun ö ventricular esquerda [New echocardiographic parameters for left ventricular function analysis]. <i>Experts in Ultrasound Reviews and Perspectives</i> , 2010 , 2, 198-204			

58	Myocardial Perfusion Imaging with Contrast Echocardiography. 2010, 297-308		
57	Intravenous administration of nicorandil immediately before percutaneous coronary intervention can prevent slow coronary flow phenomenon. <i>Okayama Igakkai Zasshi</i> , 2010 , 122, 101-106	O	
56	Redefining the Success of Mechanical Reperfusion. 2010 , 221-226		
55	Redefining the Success of Mechanical Reperfusion. 2010 , 227-233		
54	Redefining the Success of Mechanical Reperfusion. 2010 , 203-214		
53	Angiographic impact of the GuardWire system on inflated coronary segments after six months: does the distal protection balloon of the GuardWire Plusâllead to restenosis?. <i>EuroIntervention</i> , 2010 , 6, 257-260	3.1	
52	Myocardial Contrast Echocardiography After Myocardial Infarction. 2011, 231-232		
51	Featured Expert Commentary and Review: Myocardial Contrast Echocardiography for Chest Pain Syndromes in the Emergency Department. 2011 , 239-243		
50	Use of Thrombectomy Devices in Primary Percutaneous Interventions for ST-elevation Myocardial Infarction - An Update. <i>Interventional Cardiology Review</i> , 2014 , 9, 102-107	4.2 1	
49	IRM de la micro-obstruction vasculaire dans lâIhfarctus du myocarde aigu. 2011 , 179-194		
48	Myocardial Contrast Echocardiography in the Evaluation of Hypertensive Heart Disease. <i>Cardiology Research</i> , 2011 , 2, 259-268	1.8 1	
47	No Reflow. 417-432		
46	Thrombectomy and Embolic Protection. 338-353		
45	Contrast Echocardiography. 2012 , 271-283		
44	Intracoronary Stenting Strategies. 2013 , 497-529		
43	Evaluation of Medical Therapies and Intravascular Devices with Quantitative Coronary Angiography. <i>Contemporary Cardiology</i> , 2014 , 41-49	0.1	
42	Myocardial contrast echocardiography in ischemic heart disease: Osaka experience. 1997 , 387-399		
41	Intravenous echocontrast for assessment of left ventricular function and perfusion. <i>Developments in Cardiovascular Medicine</i> , 1998 , 333-350		

Interventional Approaches to the Treatment of Acute Coronary Ischemic Syndromes. *Developments in Cardiovascular Medicine*, **1998**, 277-312

	III Caralovascatar Medicine, 1996, 211-312		
39	????? : ?????(?????? IV ??????????)(?62? ?????????). Journal of JCS Cardiologists, 1998 , 6, 229-236	0.1	
38	Determining microvascular obstruction and infarct size with steady-state free precession imaging cardiac MRI. <i>PLoS ONE</i> , 2015 , 10, e0119788	3.7	1
37	Assessment of Changing Patterns of Hematological Indices and Their Effects on the Success of Primary Percutaneous Reperfusion in Acute Transmural Myocardial Infarction. <i>Research in Cardiovascular Medicine</i> , 2016 , Inpress,	0.4	
36	Myocardial Viability and Imaging in the Failing Heart. 2017, 211-223		
35	Artery diameter ratio after recanalization in endovascular therapy for acute ischemic stroke: a new predictor of clinical outcomes. <i>Neuroradiology</i> , 2021 , 1	3.2	
34	EFFECT OF PRIMARY STENTING OF CORONARY ARTERIES ON CLINICAL COURSE AND REMODELING OF THE LEFT VENTRICLE IN PATIENTS WITH ACUTE ST SEGMENT ELEVATION MYOCARDIAL INFARCTION (RESULTS AFTER 12 MONTHS). Wiadomo@i Lekarskie, 2020, 73, 68-72	0.3	
33	Angiographic determination of myocardial reperfusion by primary coronary angioplasty for acute myocardial infarction. <i>Netherlands Heart Journal</i> , 2002 , 10, 111-117	2.2	3
32	Angiographic determinants of infarct size after successful percutaneous intervention for acute ST-elevation myocardial infarction: the impact of distal embolisation. <i>Netherlands Heart Journal</i> , 2002 , 10, 353-359	2.2	2
31	The diagnosis and treatment of the no-reflow phenomenon in patients with myocardial infarction undergoing percutaneous coronary intervention. <i>Experimental and Clinical Cardiology</i> , 2008 , 13, 121-8		28
30	The Importance of Amino-terminal pro-Brain Natriuretic Peptide Testing in Clinical Cardiology. <i>Biomarker Insights</i> , 2007 , 1, 143-55	3.5	
29	The relation between myocardial blush grade and myocardial contrast echocardiography: which one is a better predictor of myocardial damage?. <i>Netherlands Heart Journal</i> , 2010 , 18, 25-30	2.2	6
28	MRI manifestations of persistent microvascular obstruction and acute left ventricular remodeling in an experimental reperfused myocardial infarction. <i>Quantitative Imaging in Medicine and Surgery</i> , 2012 , 2, 12-20	3.6	7
27	Early Administration of Intracoronary Nitroprusside Compared with Thrombus Aspiration in Myocardial Perfusion for Acute Myocardial Infarction: A 3-Year Clinical Follow-Up Study. <i>Acta Cardiologica Sinica</i> , 2015 , 31, 373-80	1.1	1
26	Renal Dysfunction on Admission Predicts No-Reflow Phenomenon in Patients Undergoing Manual Thrombus Aspiration during Primary Percutaneous Coronary Intervention. <i>Acta Cardiologica Sinica</i> , 2016 , 32, 185-93	1.1	8
25	Cutoff Value of Admission N-Terminal Pro-Brain Natriuretic Peptide Which Predicts Poor Myocardial Perfusion after Primary Percutaneous Coronary Intervention for ST-Segment-Elevation Myocardial Infarction. <i>Acta Cardiologica Sinica</i> , 2016 , 32, 649-655	1.1	2
24	Intracoronary hypothermia for acute myocardial infarction in the isolated beating pig heart. <i>American Journal of Translational Research (discontinued)</i> , 2017 , 9, 558-568	3	8
23	Early administration of cold water and adipose derived mesenchymal stem cell derived exosome effectively protects the heart from ischemia-reperfusion injury. <i>American Journal of Translational Research (discontinued)</i> , 2019 , 11, 5375-5389	3	5

22	Intra-Coronary Administration of Tacrolimus Improves Myocardial Perfusion and Left Ventricular Function in Patients with ST-Segment Elevation Myocardial Infarction (COAT-STEMI) Undergoing Primary Percutaneous Coronary Intervention. <i>Acta Cardiologica Sinica</i> , 2021 , 37, 239-253	1.1	
21	No-reflow phenomenon and in vivo cholesterol crystals combined with lipid core in acute myocardial infarction <i>IJC Heart and Vasculature</i> , 2022 , 38, 100953	2.4	O
20	Intracoronary epinephrine and verapamil in the refractory no-reflow phenomenon in patients with acute myocardial infarction. <i>Cardiovascular Therapy and Prevention (Russian Federation)</i> , 2022 , 21, 2936	0.9	
19	Modulation of cardiac remodeling by adenosine: in vitro and in vivo effects. <i>Molecular and Cellular Biochemistry</i> , 2003 , 251, 17-26	4.2	12
18	Association between cine CMR-based radiomics signature and microvascular obstruction in patients with ST-segment elevation myocardial infarction <i>Journal of Thoracic Disease</i> , 2022 , 14, 969-978	2.6	0
17	Prophylactic use of intracoronary sodium nitroprusside decreases incidence of no reflow after primary angioplasty in acute myocardial infarction. 2021 , 10, 1		
16	Comparison of angiographic results and clinical outcomes of no-reflow after stenting in left anterior descending (LAD) versus non-LAD culprit STEMI SAGE Open Medicine, 2022, 10, 205031212210	088 ¹ 106	5
15	A Narrative Review of the Classical and Modern Diagnostic Methods of the No-Reflow Phenomenon <i>Diagnostics</i> , 2022 , 12,	3.8	O
14	Mesenchymal Stem Cell-Originated Exosomal Lnc A2M-AS1 Alleviates Hypoxia/Reperfusion-Induced Apoptosis and Oxidative Stress in Cardiomyocytes <i>Cardiovascular Drugs and Therapy</i> , 2022 ,	3.9	1
13	Assessment and Treatment for Coronary Microvascular Dysfunction by Contrast Enhanced Ultrasound. <i>Frontiers in Cardiovascular Medicine</i> , 9,	5.4	1
12	Combined therapy with dapagliflozin and entresto offers an additional benefit on improving the heart function in rat after ischemia-reperfusion injury. <i>Biomedical Journal</i> , 2022 ,	7.1	
11	Efficacy of Shexiang Tongxin Dropping Pills in a Swine Model of Coronary Slow Flow. <i>Frontiers in Physiology</i> , 13,	4.6	O
10	Use of a Cutting Balloon Reduces the Incidence of Distal Embolism in Acute Coronary Syndrome Requiring Predilatation Before Stenting. <i>Circulation Reports</i> , 2022 ,	0.7	
9	Prognostic significance of thrombus burden on short- and long-term clinical outcomes in patients with ST-elevation myocardial infarction undergoing primary percutaneous coronary intervention. Publish Ahead of Print,		
8	Multicenter clinical trials that take cardiovascular care one step further. 2022 , 134, 79-85		O
7	Multimodality Imaging of Sudden Cardiac Death and Acute Complications in Acute Coronary Syndrome. 2022 , 11, 5663		O
6	Soluble ST2 in coronary artery disease: Clinical biomarkers and treatment guidance. 9,		O
5	Avalia® do N¤el de Fluxo Coron®io com MOTS-C em Pacientes com IAMCSST Submetidos ÎCP Prim®ia. 2023 , 120,		O

CITATION REPORT

Low Quantitative Blush Evaluator score predicts larger infarct size and reduced left ventricular systolic function in patients with STEMI regardless of diabetes status. 2023, 13,

Dynamic Behavior of Polymer Microbubbles During Long Ultrasound Tone-Burst Excitation and Its Application for Sonoreperfusion Therapy. 2023,

Identification of key genes and lncRNAs based on M6A methylation regulators and Ferroptosis related genes for the slow flow/no reflow phenomenon.

Sustained Mechanical Aspiration Thrombectomy for High Thrombus Burden Coronary Vessel Occlusion: The Multicenter CHEETAH Study. 2023, 16,