CITATION REPORT List of articles citing

Exercise myocardial perfusion SPECT in patients without known coronary artery disease: incremental prognostic value and use in risk stratification

DOI: 10.1161/01.CIR.93.5.905 Circulation, 1996, 93, 905-14.

Source: https://exaly.com/paper-pdf/86224892/citation-report.pdf

Version: 2024-04-11

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
673	Identification of severe and extensive coronary artery disease by automatic measurement of transient ischemic dilation of the left ventricle in dual-isotope myocardial perfusion SPECT. 1996 , 27, 1612-20		195
672	Impact of myocardial perfusion single-photon emission computed tomography on referral to catheterization of the very elderly. Is there evidence of gender-related referral bias?. 1996 , 28, 680-6		35
671	Risk assessment in patients with stable coronary artery disease: incremental value of nuclear imaging. <i>Journal of Nuclear Cardiology</i> , 1996 , 3, S41-9	2.1	16
670	A primer of biostatistic and economic methods for diagnostic and prognostic modeling in nuclear cardiology: Part I. <i>Journal of Nuclear Cardiology</i> , 1996 , 3, 538-45	2.1	14
669	Relation of gender to physician use of test results and to the prognostic value of stress technetium 99m sestamibi myocardial single-photon emission computed tomography scintigraphy. 1997 , 134, 73-8	2	44
668	Evaluation of exercise thallium scintigraphy versus exercise electrocardiography in predicting survival outcomes and morbid cardiac events in patients with single- and double-vessel disease. Findings from the Angioplasty Compared to Medicine (ACME) Study. 1997 , 30, 1256-63		59
667	Value of gating of technetium-99m sestamibi single-photon emission computed tomographic imaging. 1997 , 30, 1687-92		186
666	Cardiac function, perfusion, and morbidity in irradiated long-term survivors of Hodgkin's disease. 1997 , 39, 897-906		77
665	Can single photon emission computed tomography myocardial perfusion imaging monitor the potential benefit of aggressive treatment of hyperlipidemia?. <i>Journal of Nuclear Cardiology</i> , 1997 , 4, 555-68	2.1	6
664	Risk-sensitive therapeutic strategies for coronary artery disease: toward testing-driven therapy in stable angina patients with low-to-intermediate risk cardiac imaging results. <i>Journal of Nuclear Cardiology</i> , 1997 , 4, 409-17	2.1	3
663	Usefulness of hemodynamic changes during adenosine infusion in predicting the diagnostic accuracy of adenosine technetium-99m sestamibi single-photon emission computed tomography (SPECT). 1997 , 79, 1319-22		33
662	Identification of severe or extensive coronary artery disease in women by adenosine technetium-99m sestamibi SPECT. 1997 , 80, 132-7		93
661	Risk stratification in coronary artery disease: implications for stabilization and prevention. 1997 , 79, 10	-6	65
660	Achieving sustained improvement in myocardial perfusion: role of isosorbide mononitrate. 1997 , 79, 31-5		9
659	Incremental prognostic value of adenosine stress myocardial perfusion single-photon emission computed tomography and impact on subsequent management in patients with or suspected of having myocardial ischemia. 1997 , 80, 426-33		106
658	Predicting adverse outcome with exercise SPECT technetium-99m sestamibi imaging in patients with suspected or known coronary artery disease. 1997 , 79, 270-4		46
657	Potential cost effectiveness of initial myocardial perfusion imaging for assessment of emergency department patients with chest pain. 1997 , 79, 595-9		79

656	Impact of contrast echocardiography on diagnostic algorithms: pharmacoeconomic implications. 1997 , 20, 139-48		33
655	Treadmill test scores: attributes and limitations. <i>Journal of Nuclear Cardiology</i> , 1997 , 4, 74-8	2.1	13
654	Prediction of severe coronary artery disease by combined rest and exercise radionuclide angiocardiography and tomographic perfusion imaging with technetium 99m-labeled sestamibi: a comparison with clinical and electrocardiographic data. <i>Journal of Nuclear Cardiology</i> , 1997 , 4, 189-94	2.1	18
653	Attenuation-corrected rest thallium-201/stress technetium 99m sestamibi myocardial SPECT in normals. <i>Journal of Nuclear Cardiology</i> , 1998 , 5, 48-55	2.1	22
652	Prognostic value of stress echocardiography in the evaluation of atypical chest pain patients without known coronary artery disease. 1998 , 81, 545-51		34
651	Impact of exercise single-photon emission computed tomographic imaging on appropriateness of coronary revascularization. 1998 , 81, 1489-91		7
650	Improved myocardial ischemia detection by combined physical and mental stress testing. 1998 , 82, 109-	13	9
649	Comparison of post-stress ejection fraction and relative left ventricular volumes by automatic analysis of gated myocardial perfusion single-photon emission computed tomography acquired in the supine and prone positions. <i>Journal of Nuclear Cardiology</i> , 1998 , 5, 40-7	2.1	26
648	Prognostic characterization of patients with mild coronary artery disease with myocardial perfusion single photon emission computed tomography: validation of an outcomes-based strategy. <i>Journal of Nuclear Cardiology</i> , 1998 , 5, 90-5	2.1	3
647	Thallium perfusion defects and impaired exercise capacity predict death and non-fatal MI in symptom-free patients with previous CABG. 1998 , 2, 73		
646	Prognostic value of exercise thallium-201 imaging performed within 2 years of coronary artery bypass graft surgery. 1998 , 31, 848-54		50
645	Risk assessment using single-photon emission computed tomographic technetium-99m sestamibi imaging. 1998 , 32, 57-62		343
644	Comparative prognostic value of automatic quantitative analysis versus semiquantitative visual analysis of exercise myocardial perfusion single-photon emission computed tomography. 1998 , 32, 1987	'-95	117
643	Risk stratification with pre-hospital discharge exercise technetium-99m sestamibi myocardial tomography in men after acute myocardial infarction. 1998 , 136, 87-93		12
642	Advantages of diagnostic nuclear medicine part 2: cardiac and other nonmusculoskeletal disorders. 1999 , 27, 51-7		
641	The challenge of quantifying defect size and severity: reality versus algorithm. <i>Journal of Nuclear Cardiology</i> , 1999 , 6, 362-71	2.1	6
640	Quantification of SPECT myocardial perfusion images: methodology and validation of the Yale-CQ method. <i>Journal of Nuclear Cardiology</i> , 1999 , 6, 190-204	2.1	51
639	Using an outcomes-based approach to identify candidates for risk stratification after exercise treadmill testing. 1999 , 14, 1-9		23

638	The prognostic value of a normal Tc-99m sestamibi SPECT study in suspected coronary artery disease. <i>Journal of Nuclear Cardiology</i> , 1999 , 6, 252-6	2.1	35
637	Advances in Exercise Echocardiography Can This Technique Still Thrive in the Era of Pharmacologic Stress Testing?. 1999 , 16, 841-856		4
636	Segmental myocardial contractility versus perfusion in Kawasaki disease with coronary arterial aneurysm. 1999 , 83, 48-51		12
635	Comparison of the polar maps method and the summed stress score for predicting outcome in medically treated patients with coronary artery disease. 1999 , 83, 258-9, A5		2
634	Comparison of the prognostic value of qualitative versus quantitative stress tomographic perfusion imaging. 1999 , 83, 945-8, A9		6
633	Usefulness of an exaggerated systolic blood pressure response to exercise in predicting myocardial perfusion defects in known or suspected coronary artery disease. 1999 , 84, 1304-10		24
632	Long-term prognostic value of Duke treadmill score and exercise thallium-201 imaging performed one to three years after percutaneous transluminal coronary angioplasty. 1999 , 84, 1323-7		26
631	The ongoing evolution of risk stratification using myocardial perfusion imaging in patients with known or suspected coronary artery disease. 1999 , 8, 66-71		6
630	The economic consequences of available diagnostic and prognostic strategies for the evaluation of stable angina patients: an observational assessment of the value of precatheterization ischemia. Economics of Noninvasive Diagnosis (END) Multicenter Study Group. 1999 , 33, 661-9		282
629	Value of negative dobutamine stress echocardiography in predicting long-term cardiac events. 1999 , 12, 471-5		15
628	Screening for coronary artery disease. 1999 , 83, 1375-402, v-vi		6
627	ACC/AHA/ACP-ASIM guidelines for the management of patients with chronic stable angina: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee on Management of Patients With Chronic Stable Angina). 1999 , 33, 2092-197		482
626	The role of nuclear cardiology in clinical decision making. 1999 , 29, 280-97		38
625	Pharmacological stress testing. 1999 , 29, 298-318		49
624	Stress myocardial perfusion imaging versus echocardiography for the diagnosis and risk stratification of patients with known or suspected coronary artery disease. 1999 , 29, 319-29		4
623	Incremental prognostic value of myocardial perfusion single photon emission computed tomography in patients with diabetes mellitus. 1999 , 138, 1025-32		143
622	Comparative ability of myocardial perfusion single-photon emission computed tomography to detect coronary artery disease in patients with and without diabetes mellitus. 1999 , 137, 949-57		109
621	Technetium Tc 99m sestamibi myocardial perfusion imaging: current role for evaluation of prognosis. 1999 , 115, 1684-94		21

620 Silent myocardial ischemia and low aerobic capacity: an unlucky combination. 1999, 47, 1026-8

619	Exercise-induced silent myocardial ischemia and future cardiac events in healthy, sedentary, middle-aged and older men. 1999 , 47, 923-9		16
618	Comparative localization of myocardial ischemia by exercise electrocardiography and myocardial perfusion SPECT. <i>Journal of Nuclear Cardiology</i> , 2000 , 7, 140-5	2.1	8
617	Palpable cardiac impulse predicts adequate acoustic windows. 2000 , 17, 1-6		1
616	Exercise QT Dispersion as an Independent Predictor of the Presence of Ischemia on Myocardial Perfusion Imaging. 2000 , 5, 240-247		
615	The ongoing evolution of risk stratification using myocardial perfusion imaging in patients with known or suspected coronary artery disease. 2000 , 9, S9-S14		
614	Cost-effectiveness of stress echocardiography and nuclear perfusion imaging. 2000 , 43, 197-214		1
613	Methodological considerations in the assessment of noninvasive testing using outcomes research: pitfalls and limitations. 2000 , 43, 215-30		20
612	Stress nuclear myocardial perfusion imaging versus stress echocardiography: prognostic comparisons. 2000 , 43, 231-44		5
611	Evaluation of chest pain in patients with low to intermediate pretest probability of coronary artery disease by electron beam computed tomography. 2000 , 85, 283-8		42
610	Noninvasive strategies for the estimation of cardiac risk in stable chest pain patients. The Economics of Noninvasive Diagnosis (END) Study Group. 2000 , 86, 1-7		61
609	Comparison of baseline and low-dose dobutamine technetium-99m sestamibi scintigraphy with low-dose dobutamine echocardiography for predicting functional recovery after revascularization. 2000 , 86, 153-7		13
608	Sustained reduction of exercise perfusion defect extent and severity with isosorbide mononitrate (Imdur) as demonstrated by means of technetium 99m sestamibi. <i>Journal of Nuclear Cardiology</i> , 2000 , 7, 342-53	2.1	21
607	Are the American College of Cardiology/American Heart Association guidelines for exercise testing for suspected coronary artery disease correct?. 2000 , 118, 535-41		17
606	Myocardial perfusion imaging. 2000 , 83, 355-60		26
605	Focal angiogen therapy using intramyocardial delivery of an adenovirus vector coding for vascular endothelial growth factor 121. 2000 , 69, 14-23; discussion 23-4		120
604	Patients with intermediate-risk exercise tests but without perfusion defects and normal cardiac sizes were at low risk of cardiovascular events. 2000 , 4, 41		
603	Perfusion imaging. 2000 , 35, 30B-31B		

602	Identification of severe coronary artery disease in patients with a single abnormal coronary territory on exercise thallium-201 imaging: the importance of clinical and exercise variables. 2000 , 35, 335-44		24
601	When to stress patients after coronary artery bypass surgery? Risk stratification in patients early and late post-CABG using stress myocardial perfusion SPECT: implications of appropriate clinical strategies. 2001 , 37, 144-52		93
600	Prediction of mortality using dobutamine echocardiography. 2001 , 37, 754-60		146
599	Gender differences and temporal trends in clinical characteristics, stress test results and use of invasive procedures in patients undergoing evaluation for coronary artery disease. 2001 , 38, 690-7		65
598	Peripheral vascular endothelial function testing as a noninvasive indicator of coronary artery disease. 2001 , 38, 1843-9		204
597	Exercise electrocardiogram testing and prognosis. Novel markers and predictive instruments. 2001 , 19, 401-14		36
596	Assessing patients with possible heart disease using scores. 2001 , 31, 387-408		1
595	Medical therapy for coronary artery disease works, even (especially) in the real world. 2001 , 110, 497-8		6
594	Cardiac nuclear medicine in monitoring patients with coronary heart disease. 2001 , 31, 223-37		21
593	[Angiographic correlates of the high-risk criteria for conventional exercise testing and the Duke treadmill score]. 2001 , 54, 860-7		2
592	Thallium gated SPECT: relation between immediate post-stress evolution of ejection fraction and severity of perfusion pattern. 2001 , 22, 57-64		5
591	Interpretive intra- and interobserver reproducibility of rest/stress 99Tcm-sestamibi myocardial perfusion SPECT in a consecutive group of male patients with stable angina pectoris before and after percutaneous transluminal angioplasty. 2001 , 22, 531-7		25
590	Preliminary clinical results of photon energy recovery in simultaneous rest Tl-201/stress Tc-99m sestamibi myocardial SPECT. <i>Journal of Nuclear Cardiology</i> , 2001 , 8, 144-51	2.1	4
589	Post-myocardial infarction risk stratification with stress nuclear myocardial perfusion imaging versus echocardiography: separate but not equal. <i>Journal of Nuclear Cardiology</i> , 2001 , 8, 215-8	2.1	1
588	Serial changes on quantitative myocardial perfusion SPECT in patients undergoing revascularization or conservative therapy. <i>Journal of Nuclear Cardiology</i> , 2001 , 8, 428-37	2.1	25
587	What is wrong with the treadmill exercise test?. <i>Journal of Nuclear Cardiology</i> , 2001 , 8, 415-6	2.1	
586	Prediction of severe coronary artery disease and long-term outcome in patients undergoing vasodilator SPECT. <i>Journal of Nuclear Cardiology</i> , 2001 , 8, 438-44	2.1	18
585	Incidence of major cardiovascular events in black patients with normal myocardial stress perfusion study results. <i>Journal of Nuclear Cardiology</i> , 2001 , 8, 541-7	2.1	12

(2002-2001)

584	Beyond the cholesterol profile: monitoring therapeutic effectiveness of statin therapy. <i>Journal of Nuclear Cardiology</i> , 2001 , 8, 528-32	5
583	Racial disparities in cardiovascular health care: unfulfilled exSPECTations?. <i>Journal of Nuclear Cardiology</i> , 2001 , 8, 626-9	2
582	Usefulness of worsening clinical status or exercise performance in predicting future events in patients with coronary artery disease. 2001 , 88, 1294-7	5
581	Principal uses of myocardial perfusion scintigraphy in the management of patients with known or suspected coronary artery disease. 2001 , 43, 281-302	22
580	Recent advances in myocardial perfusion imaging. 2001 , 26, 1-140	4
579	Stress testing in cardiac evaluation: current concepts with emphasis on the ECG. 2001, 119, 907-25	41
578	Clinical practice. Noninvasive tests in patients with stable coronary artery disease. 2001, 344, 1840-5	118
577	Prognostic value of dobutamine-atropine stress myocardial perfusion imaging in patients with diabetes. 2002 , 25, 1637-43	19
576	Exercise testing scores as an example of better decisions through science. 2002 , 34, 1391-8	15
575	Validation of quantitative gated single photon emission computed tomography and an automated scoring system for the assessment of regional left ventricular systolic function. 2002 , 23, 887-98	19
574	High coronary artery calcium scores pose an extremely elevated risk for hard events. 2002 , 39, 225-30	249
573	Coronary flow studies for risk stratification in multivessel disease. A physiologic bridge too far?. 2002 , 39, 859-63	7
57 ²	Reversible regional wall motion abnormalities on exercise technetium-99m-gated cardiac single photon emission computed tomography predict high-grade angiographic stenoses. 2002 , 39, 991-8	93
571	Low event rate for stress-only perfusion imaging in patients evaluated for chest pain. 2002 , 39, 999-1004	81
57°	The "exercise" part of exercise echocardiography. 2002 , 39, 1353-5	8
569	ACC/AHA 2002 guideline update for exercise testing: summary article. A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee to Update the 1997 Exercise Testing Guidelines). 2002 , 40, 1531-40	976
568	Use of a simple clinical score to predict prognosis of patients with normal or mildly abnormal resting electrocardiographic findings undergoing evaluation for coronary artery disease. 2002 , 77, 515-21	35
567	Exercise testing for assessment of autonomic function. 2002 , 144, 580-2	1

566	Exercise echocardiography and thallium-201 single-photon emission computed tomography stress test for 5- and 10-year prognosis of mortality and specific cardiac events. 2002 , 15, 1326-34		13
565	Las pruebas de esfuerzo. 2002 , 14, 41-50		
564	Stress echocardiography for risk stratification of patients with chest pain and normal or slightly narrowed coronary arteries. 2002 , 15, 1285-9		9
563	Better decisions through science: exercise testing scores. 2002 , 44, 395-414		14
562	Prevalence and prognostic value of perfusion defects detected by stress technetium-99m sestamibi myocardial perfusion single-photon emission computed tomography in asymptomatic patients with diabetes mellitus and no known coronary artery disease. 2002 , 90, 827-32		130
561	Atypical chest pain, myocardial perfusion SPET and left main stem occlusion. 2002 , 29, 718		
560	Risk stratification in patients with remote prior myocardial infarction using rest-stress myocardial perfusion SPECT: prognostic value and impact on referral to early catheterization. <i>Journal of Nuclear Cardiology</i> , 2002 , 9, 23-32	2.1	39
559	Cardiac risk defined by stress myocardial perfusion imaging: Impact on physician decision making and cost savings. <i>Journal of Nuclear Cardiology</i> , 2002 , 9, 124-6	2.1	
558	Comparison of the cost-effectiveness of stress myocardial SPECT and stress echocardiography in suspected coronary artery disease considering the prognostic value of false-negative results. <i>Journal of Nuclear Cardiology</i> , 2002 , 9, 515-22	2.1	21
557	Exercise myocardial perfusion imaging in elderly patients. <i>Journal of Nuclear Cardiology</i> , 2002 , 9, 573-80	2.1	26
556	And what do we tell our elders?. Journal of Nuclear Cardiology, 2002, 9, 672-4	2.1	2
555	Use of atropine in patients with submaximal heart rate during exercise myocardial perfusion SPECT. <i>Journal of Nuclear Cardiology</i> , 2003 , 10, 51-5	2.1	7
554	Analysis of referral patterns, predictive accuracy, and impact on patient management of myocardial perfusion imaging in a new nuclear cardiology laboratory. <i>Journal of Nuclear Cardiology</i> , 2003 , 10, 148-5	3 ^{2.1}	9
553	Risk stratification of the normal perfusion scan: does normal stress perfusion always mean very low risk?. <i>Journal of Nuclear Cardiology</i> , 2003 , 10, 87-91	2.1	8
552	Do vascular surgery patients need a cardiology work-up? A review of pre-operative cardiac clearance guidelines in vascular surgery. 2003 , 25, 110-7		16
551	Diagnosis of coronary artery disease in women. 2003 , 46, 239-58		9
550	Better decisions through science: exercise testing scores. 2003 , 28, 589-620		1
549	[Magnetic resonance imaging early after acute myocardial infarction. A visual analysis of myocardial perfusion based on a 17 segment model]. 2003 , 52, 7-14		4

548	Prognostic value of exercise cardiac tomography performed late after percutaneous coronary intervention in symptomatic and symptom-free patients. 2003 , 91, 259-63		32
547	Incremental value of exercise technetium-99m tetrofosmin myocardial perfusion single-photon emission computed tomography for the prediction of cardiac events. 2003 , 91, 408-11		38
546	Relation of local platelet glycoprotein IIb/IIIa independent activation during coronary angioplasty in acute myocardial infarction to recovery of left ventricular function. 2003 , 92, 446-50		
545	Rest perfusion defects in patients with no history of myocardial infarction predict the presence of a critical coronary artery stenosis. <i>Journal of Nuclear Cardiology</i> , 2003 , 10, 656-62	2.1	3
544	Long-term prognosis after a normal exercise stress Tc-99m sestamibi SPECT study. <i>Journal of Nuclear Cardiology</i> , 2003 , 10, 261-6	2.1	53
543	Antianginal medications and diagnostic accuracy of myocardial perfusion imaging. <i>Journal of Nuclear Cardiology</i> , 2003 , 10, 433-5	2.1	2
542	First annual Mario S. Verani, MD, Memorial lecture: clinical value of myocardial perfusion imaging in coronary artery disease. <i>Journal of Nuclear Cardiology</i> , 2003 , 10, 529-42	2.1	22
541	Myocardial Perfusion Imaging. 2003 , 7, 29-35		
540	Esophageal dysfunction in patients with atypical chest pain investigated with esophageal scintigraphy and myocardial perfusion imaging: an outcome study. <i>Journal of Nuclear Cardiology</i> , 2003 , 10, 490-7	2.1	10
539	[Value of gated-SPECT in defining the post-revascularization prognosis of patients with ischemic cardiomyopathy]. 2003 , 56, 281-8		2
538	ACC/AHA/ASNC guidelines for the clinical use of cardiac radionuclide imagingexecutive summary: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (ACC/AHA/ASNC Committee to Revise the 1995 Guidelines for the Clinical Use of Cardiac Radionuclide Imaging). 2003, 42, 1318-33		580
537	Determinants of risk and its temporal variation in patients with normal stress myocardial perfusion scans: what is the warranty period of a normal scan?. 2003 , 41, 1329-40		288
536	34th Bethesda Conference: Task force #1Identification of coronary heart disease risk: is there a detection gap?. 2003 , 41, 1863-74		147
535	Long-term outcome of patients with silent versus symptomatic ischemia six months after percutaneous coronary intervention and stenting. 2003 , 42, 33-40		87
534	Prospective serial evaluation of myocardial perfusion and lipids during the first six months of pravastatin therapy: coronary artery disease regression single photon emission computed tomography monitoring trial. 2003 , 42, 600-10		52
533	Evaluation of pretest and exercise test scores to assess all-cause mortality in unselected patients presenting for exercise testing with symptoms of suspected coronary artery disease. 2003 , 42, 842-50		66
532	Is there a referral bias against catheterization of patients with reduced left ventricular ejection fraction? Influence of ejection fraction and inducible ischemia on post-single-photon emission computed tomography management of patients without a history of coronary artery disease. 2003 ,		31
531	42, 1286-94 Role of myocardial perfusion imaging for risk stratification in suspected or known coronary artery disease. 2003 , 89, 1291-7		49

530	Arteriogenesis induced by intramyocardial vascular endothelial growth factor 165 gene transfer in chronically ischemic pigs. 2003 , 14, 1307-18	39
529	Contemporary cardiology and hysteric nucleophobia. 2003 , 114, 131-4	9
528	[Clinical predictors of the presence of perfusion defects in more than than one territory in myocardial scintigraphy with GATED-SPECT]. 2003 , 22, 20-5	
527	Finding the age of the patient's heart. 2003 , 326, 1045-6	4
526	Procedure guidelines for radionuclide myocardial perfusion imaging. 2003 , 24, 1105-1119	18
525	New frontiers in risk stratification using stress myocardial perfusion single photon emission computed tomography. 2003 , 18, 494-502	5
524	Prediction of cardiac death in hypertensive patients with suspected or known coronary artery disease by stress technetium-99m tetrofosmin myocardial perfusion imaging. 2003 , 21, 1945-51	16
523	Stress echocardiography versus myocardial SPECT for risk stratification of patients with coronary artery disease. 2003 , 18, 486-93	8
522	Critical pathway for an aggressive diagnostic strategy for patients who present with chest pain but a nondiagnostic electrocardiogram to the emergency department. 2003 , 2, 88-94	1
521	Outcome prediction in patients at high risk for coronary artery disease: comparison between 99mTc tetrofosmin and 99mTc sestamibi. 2004 , 232, 58-65	17
520	Assessment of prognosis in chronic coronary artery disease. <i>Heart</i> , 2004 , 90 Suppl 5, v10-5 5.1	8
519	Procedure guidelines for radionuclide myocardial perfusion imaging. 2004 , 90 Suppl 1, i1-10	46
518	Prognostic value of exercise stress myocardial perfusion imaging in patients with permanent pacemakers. 2004 , 94, 811-4	7
517	Quantitative relation of ST-segment depression during exercise to the magnitude of myocardial ischemia as assessed by single-photon emission computed tomographic myocardial perfusion imaging. 2004 , 94, 703-8	14
516	Prognostic stratification of patients with right bundle branch block using dobutamine stress echocardiography. 2004 , 94, 954-7	7
515	Dobutamine stress MRI. Part II. Risk stratification with dobutamine cardiovascular magnetic resonance in patients suspected of myocardial ischemia. 2004 , 14, 2046-52	49
514	Myocardial perfusion scintigraphy: the evidence. 2004 , 31, 261-91	352
513	Quantification of myocardial perfusion defects using three different software packages. 2004 , 31, 229-32	18

(2004-2004)

512	Quantitative Tc-99m sestamibi attenuation-corrected SPECT: development and multicenter trial validation of myocardial perfusion stress gender-independent normal database in an obese population. <i>Journal of Nuclear Cardiology</i> , 2004 , 11, 263-72	2.1	81
511	In retroSPECT: the prognostic value of nuclear cardiology-past, present, and future. <i>Journal of Nuclear Cardiology</i> , 2004 , 11, 237-8	2.1	
510	Prognostic value of stress Tc-99m tetrofosmin SPECT in patients with previous myocardial infarction: impact of scintigraphic extent of coronary artery disease. <i>Journal of Nuclear Cardiology</i> , 2004 , 11, 704-9	2.1	17
509	Prognostic value of gated myocardial perfusion SPECT. Journal of Nuclear Cardiology, 2004, 11, 171-85	2.1	438
508	Clinical value of attenuation correction in stress-only Tc-99m sestamibi SPECT imaging. <i>Journal of Nuclear Cardiology</i> , 2004 , 11, 273-81	2.1	89
507	Prognostic validation of a 17-segment score derived from a 20-segment score for myocardial perfusion SPECT interpretation. <i>Journal of Nuclear Cardiology</i> , 2004 , 11, 414-23	2.1	196
506	Observer variability in the evaluation of dual-isotope Tl-201/Tc-99m sestamibi rest/stress myocardial perfusion SPECT in men and women with known or suspected stable angina pectoris. <i>Journal of Nuclear Cardiology</i> , 2004 , 11, 710-8	2.1	22
505	Stress myocardial perfusion single-photon emission computed tomography is clinically effective and cost effective in risk stratification of patients with a high likelihood of coronary artery disease (CAD) but no known CAD. 2004 , 43, 200-8		167
504	Utility of myocardial perfusion imaging in patients with low-risk treadmill scores. 2004 , 43, 194-9		61
503	Risk stratification using stress myocardial perfusion imaging: don't neglect the value of clinical variables. 2004 , 43, 209-12		8
502	Relationship between stress-induced myocardial ischemia and atherosclerosis measured by coronary calcium tomography. 2004 , 44, 923-30		339
501	A novel method for the detection of transient myocardial ischaemia using body surface electrocardiac mapping. 2004 , 95, 75-81		10
500	Yield of stress single-photon emission computed tomography in asymptomatic patients with diabetes. 2004 , 147, 890-6		79
499	Are exercise stress tests appropriate for assessing myocardial ischemia in patients with major depressive disorder?. 2004 , 148, 621-7		43
498	Long-term prognostic value of dobutamine stress echocardiography compared with myocardial perfusion scanning in patients unable to perform exercise tests. 2004 , 117, 1-9		40
497	Role of nuclear cardiology in advancing cardiac surgery. 2004 , 16, 255-65		1
496	Incremental prognostic value of myocardial perfusion 99m-technetium-sestamibi SPECT in the elderly. 2004 , 93, 137-43		29
495	Metabolic syndrome and ischemic heart disease in gout. 2004 , 10, 105-9		41

494	Automated quantification of 99mTc sestamibi myocardial perfusion compared with visual analysis. 2004 , 25, 833-8		6
493	The prognostic value of myocardial perfusion imaging in patients with suspected or known coronary artery disease. 2004 , 25, 217-20		1
492	Diabetes mellitus: evaluation of patients with known or suspected coronary artery disease and the role played by myocardial perfusion imaging. 2005 , 26, 587-91		2
491	Radionuclide-based insights into the pathophysiology of ischemic heart disease: beyond diagnosis. 2005 , 53, 176-91		5
490	Prognostic value of I-123 metaiodobenzylguanidine imaging in vasospastic angina without significant coronary stenosis. 2005 , 69, 171-6		24
489	Incremental value of regional wall motion analysis immediately after exercise for the detection of single-vessel coronary artery disease: study by separate acquisition, dual-isotope ECG-gated single-photon emission computed tomography. 2005 , 69, 301-5		32
488	Prediction of cardiac events in patients with transient left ventricle dilation on stress myocardial perfusion SPECT images. 2005 , 69, 1223-9		9
487	Prognostic value of myocardial perfusion SPECT versus exercise electrocardiography in patients with ST-segment depression on resting electrocardiography. <i>Journal of Nuclear Cardiology</i> , 2005 , 12, 655-61	2.1	8
486	The incidence and functional consequences of RT-associated cardiac perfusion defects. 2005 , 63, 214-2	3	352
485	The prognostic value of pharmacologic stress myocardial perfusion imaging in patients with permanent pacemakers. <i>Journal of Nuclear Cardiology</i> , 2005 , 12, 37-42	2.1	8
484	Can the reversible regional wall motion abnormalities on stress gated Tc-99m sestamibi SPECT predict a future cardiac event?. <i>Journal of Nuclear Cardiology</i> , 2005 , 12, 20-31	2.1	24
483	One good turn deserves another. <i>Journal of Nuclear Cardiology</i> , 2005 , 12, 3-4	2.1	
482	Role of regional myocardial dysfunction by gated myocardial perfusion SPECT in the prognostic evaluation of patients with coronary artery disease. <i>Journal of Nuclear Cardiology</i> , 2005 , 12, 5-8	2.1	14
481	Pharmacokinetics and safety of binodenoson after intravenous dose escalation in healthy volunteers. <i>Journal of Nuclear Cardiology</i> , 2005 , 12, 166-71	2.1	11
480	Diagnostic accuracy of myocardial perfusion imaging in a study population without post-test referral bias. <i>Journal of Nuclear Cardiology</i> , 2005 , 12, 530-7	2.1	28
479	Prognostic value of stress myocardial perfusion imaging in octogenarian population. <i>Journal of Nuclear Cardiology</i> , 2005 , 12, 671-5	2.1	15
478	Incremental prognostic value of combined perfusion and function assessment during myocardial gated SPECT in patients aged 75 years or older. <i>Journal of Nuclear Cardiology</i> , 2005 , 12, 662-70	2.1	28
477	Can the ST segment be saved?. <i>Journal of Nuclear Cardiology</i> , 2005 , 12, 622-4	2.1	1

(2005-2005)

476	Relationship of transient ischemic dilation in dipyridamole myocardial perfusion imaging and stress-induced changes of functional parameters evaluated by Tl-201 gated SPECT. <i>Journal of 2.1 Nuclear Cardiology</i> , 2005 , 12, 268-75	43
475	Use of pro-atrial natriuretic peptide in the detection of myocardial ischaemia. 2005 , 35, 450-6	3
474	Incremental prognostic power of single-photon emission computed tomographic myocardial perfusion imaging in patients with known or suspected coronary artery disease. 2005 , 95, 182-8	31
473	Comparison of mental stress-induced myocardial ischemia in coronary artery disease patients with versus without left ventricular dysfunction. 2005 , 95, 322-6	29
472	Comparison of all-cause mortality in women with known or suspected coronary artery disease referred for dobutamine stress echocardiography with normal versus abnormal test results. 2005 , 95, 1072-5	5
471	Prognostic value of combined perfusion and function by stress technetium-99m sestamibi gated SPECT myocardial perfusion imaging in patients with suspected or known coronary artery disease. 2005 , 95, 1351-7	32
470	Comparison of prognostic value of negative dobutamine stress echocardiography versus single-photon emission computed tomography after acute myocardial infarction. 2005 , 96, 13-6	5
469	Sequential single-photon emission computed tomography myocardial perfusion imaging. 2005 , 96, 28J-39J	5
468	Relation of brachial artery flow-mediated vasodilation to significant coronary artery disease in patients with peripheral arterial disease. 2005 , 96, 1337-41	42
467	Anatomic distribution of myocardial ischemia as a determinant of exercise-induced ST-segment depression. 2005 , 96, 1356-60	12
466	Usefulness of fractional flow reserve in determining the indication of target lesion revascularization. 2005 , 65, 355-60	14
465	Incremental prognostic value of dobutamine-atropine stress 99mTc-tetrofosmin myocardial perfusion imaging for predicting outcome in diabetic patients with limited exercise capacity. 2005 , 32, 1057-63	6
464	Single-photon emission computed tomography myocardial perfusion imaging in patients with diabetes. <i>Current Cardiology Reports</i> , 2005 , 7, 117-23	4
463	Long-term prediction of mortality in elderly persons by dobutamine stress echocardiography. 2005 , 60, 1333-8	27
462	The association between left ventricle diastolic dysfunction and endothelial dysfunction and the results of stress myocardial SPECT in asymptomatic patients with type 2 diabetes. 2005 , 33, 473-82	12
461	Long term outcome in patients with silent versus symptomatic ischaemia during dobutamine stress echocardiography. <i>Heart</i> , 2005 , 91, 737-42	22
460	Prognosis of patients with good exercise capacity and mildly abnormal exercise echocardiography results: identification of an at-risk subgroup. 2005 , 18, 644-8	4
459	Cardiac single-photon emission-computed tomography using combined cone-beam/fan-beam collimation. 2005 , 52, 143-153	5

458	Marked differences between patients referred for stress echocardiography and myocardial perfusion imaging studies. 2005 , 149, 888-93	6
457	Pacemaker stress echocardiography predicts cardiac events in patients with permanent pacemaker. 2005 , 118, 1381-6	16
456	Use of N-terminal pro-B-type natriuretic peptide to detect myocardial ischemia. 2005 , 118, 1287	45
455	Ecocardiografà con dobutamina en pacientes con angina crflica estable y ergometrà de riesgo bajo o intermedio: utilidad en la valoracifi del pronfitico a largo plazo*. 2005 , 58, 916-923	О
454	Dobutamine Stress Echocardiography in Patients With Stable Chronic Angina and a Low-or Medium-Risk on Exercise Testing: Usefulness for Assessing Long-Term Prognosis. 2005 , 58, 916-923	
453	The use of nuclear cardiology in clinical decision making. 2005 , 35, 62-72	53
452	Identifying high-risk asymptomatic diabetic patients who are candidates for screening stress single-photon emission computed tomography imaging. 2005 , 45, 43-9	171
451	A prognostic score for prediction of cardiac mortality risk after adenosine stress myocardial perfusion scintigraphy. 2005 , 45, 722-9	80
450	Quantitative instruments for predicting risk and benefit. 2005 , 45, 730-2	2
449	Prognostic value of myocardial perfusion contrast echocardiography in patients with suggested or known ischemic heart disease. 2006 , 19, 1203-10	8
448	Tibolone improves myocardial perfusion in postmenopausal women with ischemic heart disease: an open-label exploratory pilot study. 2006 , 47, 559-64	6
447	Mental stress provokes ischemia in coronary artery disease subjects without exercise- or adenosine-induced ischemia. 2006 , 47, 987-91	67
446	Impact of body mass index on cardiac mortality in patients with known or suspected coronary artery disease undergoing myocardial perfusion single-photon emission computed tomography. 2006 , 47, 1418-26	36
445	What is the prognostic value of myocardial perfusion imaging using rubidium-82 positron emission tomography?. 2006 , 48, 1029-39	291
444	Prognostic utility of single-photon emission computed tomography in adult patients with hypertrophic cardiomyopathy. 2006 , 151, 426-35	34
443	Use of B-type natriuretic peptide in the detection of myocardial ischemia. 2006 , 151, 1223-30	70
442	The value of stress single photon emission computed tomography in patients without known coronary artery disease presenting with dyspnea. 2006 , 152, 551-7	9
441	QT dispersion in comparison to Tl-201-SPECT for detection of myocardial ischaemia. 2006 , 113, 327-31	2

440	Evaluation of no-reflow phenomenon using 201TlCl/123I-BMIPP dual-isotope myocardial SPECT. 2006 , 73, 258-64		3	
439	Nuclear cardiology: myocardial perfusion imaging with SPECT and PET. 2006 , 18, 166-177		6	
438	Study of the predictors and lesion characteristics of ischemic heart disease patients with false negative results in stress myocardial perfusion single-photon emission tomography. 2006 , 70, 297-303		11	
437	Quantitative Analysis in Myocardial SPECT Imaging. 2006 , 471-493		3	
436	Noninvasive preclinical assessment of CAD. 2006 , 1, 309-320			
435	Prognostic utility of sestamibi lung uptake does not require adjustment for stress-related variables: a retrospective cohort study. 2006 , 6, 2		5	
434	Prognostic value of myocardial perfusion imaging in patients with known or suspected stable angina pectoris: evaluation in a setting in which myocardial perfusion imaging did not influence the choice of treatment. 2006 , 26, 288-95		10	
433	Plasmid-mediated VEGF gene transfer induces cardiomyogenesis and reduces myocardial infarct size in sheep. 2006 , 13, 1133-42		67	
432	Revascularize only for ischemia, especially if left ventricular function is poor. <i>Journal of Nuclear Cardiology</i> , 2006 , 13, 742-6	2.1		
431	A comparison of Tl-201, Tc-99m sestamibi, and Tc-99m tetrofosmin myocardial perfusion scintigraphy in patients with mild to moderate coronary stenosis. <i>Journal of Nuclear Cardiology</i> , 2006 , 13, 488-94	2.1	11	
430	Prognostic value of poststress left ventricular volume and ejection fraction by gated myocardial perfusion SPECT in women and men: gender-related differences in normal limits and outcomes. <i>Journal of Nuclear Cardiology</i> , 2006 , 13, 495-506	2.1	92	
429	Predicting therapeutic benefit from myocardial revascularization procedures: are measurements of both resting left ventricular ejection fraction and stress-induced myocardial ischemia necessary?. <i>Journal of Nuclear Cardiology</i> , 2006 , 13, 768-78	2.1	118	
428	Characteristics and outcome of octogenarian population referred for myocardial perfusion imaging: comparison with non-octogenarian population with reference to gender. 2006 , 29, 117-20		11	
427	MDCT is better than stress perfusion imaging for detecting CADAgainst. 2006 , 33, 356-9		1	
426	Prognostic value of stress 99mTc-tetrofosmin myocardial perfusion imaging in predicting all-cause mortality: a 6-year follow-up study. 2006 , 33, 1157-61		10	
425	Worsening of left ventricular ejection fraction induced by dipyridamole on Tl-201 gated myocardial perfusion imaging predicts significant coronary artery disease. <i>Journal of Nuclear Cardiology</i> , 2006 , 13, 225-32	2.1	24	
424	Usefulness of exercise myocardial scintigraphy in multivessel coronary disease after incomplete revascularization with coronary stenting. 2006 , 97, 207-15		14	
423	Comparison of coronary calcium and stress myocardial perfusion imaging in apparently healthy siblings of individuals with premature coronary artery disease. 2006 , 97, 328-33		34	

422	Myocardial perfusion scintigraphy: an important step between clinical assessment and coronary angiography in patients with stable chest pain. 2006 , 27, 3-4	8
421	The significance of carotid artery plaques in the detection of coronary artery disease in asymptomatic type 2 diabetic patients. 2006 , 34, 13-20	3
420	PCI for stable coronary disease. 2007 , 357, 414-5; author reply 417-8	30
419	Nuclear cardiac imaging for the assessment of coronary artery disease in the elderly. 2007 , 16, 355-62	3
418	Reverse perfusion-metabolism mismatch predicts good prognosis in patients undergoing cardiac resynchronization therapy: a pilot study. 2007 , 71, 126-31	20
417	Prognostic value of myocardial perfusion single-photon emission computed tomography for the prediction of future cardiac events in a Japanese population: a middle-term follow-up study. 2007 , 71, 1580-5	40
416	Prognostic value of non-obstructive CT low-dense coronary artery plaques detected by multislice computed tomography. 2007 , 71, 1898-903	58
415	Gated single-photon emission computed tomography detects subendocardial ischemia in hypertrophic cardiomyopathy. 2007 , 71, 256-60	10
414	Prognostic value of normal stress myocardial perfusion imaging in Japanese population. 2008, 72, 611-7	42
413	Prediction of mortality in patients with coronary artery disease undergoing vasodilator stress testing: a comparison between 99mTc-tetrofosmin and 99mTc-sestamibi. 2007 , 28, 457-63	1
412	The long prognostic value of wall motion abnormalities during the recovery phase of dobutamine stress echocardiography after receiving acute beta-blockade. 2007 , 18, 187-92	6
411	Relationship and prognostic value of coronary artery calcification by electron beam computed tomography to stress-induced ischemia by single photon emission computed tomography. 2007 , 153, 807-14	50
410	Effect of exercise training on ventricular function, dyssynchrony, resting myocardial perfusion, and clinical outcomes in patients with heart failure: a nuclear ancillary study of Heart Failure and A Controlled Trial Investigating Outcomes of Exercise TraiNing (HF-ACTION); design and rationale.	13
409	Prevalence and prognosis of left ventricular systolic dysfunction in asymptomatic diabetic patients without known coronary artery disease referred for stress single-photon emission computed tomography and assessment of left ventricular function. 2007 , 154, 567-74	24
408	Valuation de la faisabilit`de la tomoscintigraphie myocardique prcoce au sestamibi synchronise ^lâlectrocardiogramme. 2007 , 31, 647-655	1
407	Neurohormonal activation and left ventricular ejection fraction in patients with suspected myocardial ischemia. 2007 , 120, 248-53	6
406	Cost implications of initial computed tomography angiography as opposed to catheterization in patients with mildly abnormal or equivocal myocardial perfusion scans. 2007 , 1, 21-6	28
405	Nuclear stress testing in elderly patients: a review of its use in the assessment of cardiac risk, particularly in patients undergoing preoperative risk assessment. 2007 , 24, 467-79	4

404	The prognostic value of normal exercise myocardial perfusion imaging and exercise echocardiography: a meta-analysis. 2007 , 49, 227-37		354
403	The value of myocardial perfusion single-photon emission computed tomography in screening asymptomatic patients with atrial fibrillation for coronary artery disease. 2007 , 50, 1080-5		20
402	Mortality risk associated with ejection fraction differs across resting nuclear perfusion findings. Journal of Nuclear Cardiology, 2007 , 14, 165-73	2.1	11
401	A randomized trial of exercise treadmill ECG versus stress SPECT myocardial perfusion imaging as an initial diagnostic strategy in stable patients with chest pain and suspected CAD: cost analysis. <i>Journal of Nuclear Cardiology</i> , 2007 , 14, 174-86	2.1	35
400	A normal stress SPECT scan is an effective gatekeeper for coronary angiography. <i>Journal of Nuclear Cardiology</i> , 2007 , 14, 187-93	2.1	13
399	The price for probability: Comparing the costs of diagnostic testing strategies. <i>Journal of Nuclear Cardiology</i> , 2007 , 14, 142-4	2.1	3
398	Influence of the postexercise acquisition delay on the detection of functional abnormalities in sestamibi-gated SPECT. <i>Journal of Nuclear Cardiology</i> , 2007 , 14, 334-40	2.1	5
397	The increasing role of quantification in clinical nuclear cardiology: the Emory approach. <i>Journal of Nuclear Cardiology</i> , 2007 , 14, 420-32	2.1	141
396	Diagnostic impact of SPECT image display on assessment of obstructive coronary artery disease. Journal of Nuclear Cardiology, 2007 , 14, 659-68	2.1	8
395	American Society of Nuclear Cardiology review of the ACCF/ASNC appropriateness criteria for single-photon emission computed tomography myocardial perfusion imaging (SPECT MPI). <i>Journal of Nuclear Cardiology</i> , 2007 , 14, e26-38	2.1	43
394	Prognostic value of SPECT myocardial perfusion imaging in patients with elevated cardiac troponin I levels and atypical clinical presentation. <i>Journal of Nuclear Cardiology</i> , 2007 , 14, 53-8	2.1	13
393	Diagnostic value of left ventricular function after stress and at rest in the detection of multivessel coronary artery disease as assessed by electrocardiogram-gated SPECT. <i>Journal of Nuclear Cardiology</i> , 2007 , 14, 68-74	2.1	31
392	A hypertensive response to exercise is associated with transient ischemic dilation on myocardial perfusion SPECT imaging. <i>Journal of Nuclear Cardiology</i> , 2007 , 14, 537-43	2.1	12
391	Variation in heart rate influences the assessment of transient ischemic dilation in myocardial perfusion scintigraphy. 2007 , 7, 1		4
390	Comparison between dobutamine echocardiography and single-photon emission computed tomography for interpretive reproducibility. 2007 , 100, 1239-44		11
389	Molecular cardiovascular imaging using scintigraphic methods. 2007 , 17, 1422-32		7
388	Quantification of left ventricular volumes and ejection fraction from gated 99mTc-MIBI SPECT: validation of an elastic surface model approach in comparison to cardiac magnetic resonance imaging, 4D-MSPECT and QGS. 2007 , 34, 900-9		27
387	Safety and feasibility of atropine added in patients with sub-maximal heart rate during exercise myocardial perfusion SPECT. 2007 , 23, 511-8		3

Frlidiagnostik und Risikostratifizierung der koronaren Herzerkrankung bei mittlerer Vortestwahrscheinlichkeit durch äßatedäßPECT. **2007**, 2, V28-V36

385	The oft neglected rest study. <i>Journal of Nuclear Cardiology</i> , 2008 , 15, 739-42	2.1	3
384	Prognostic predictors and outcomes in patients with abnormal myocardial perfusion imaging and angiographically insignificant coronary artery disease. <i>Journal of Nuclear Cardiology</i> , 2008 , 15, 754-61	2.1	15
383	Comparison of SSS and SRS calculated from normal databases provided by QPS and 4D-MSPECT manufacturers and from identical institutional normals. 2008 , 35, 311-8		22
382	Differential effects comparing exercise and pharmacologic stress on left ventricular function using gated Tc-99m sestamibi SPECT. <i>Annals of Nuclear Medicine</i> , 2008 , 22, 185-90	2.5	7
381	Impact of inducible ischemia by stress SPECT in cardiac risk assessment in diabetic patients: rationale and design of a prospective, multicenter trial. <i>Journal of Nuclear Cardiology</i> , 2008 , 15, 100-4	2.1	16
380	Comparison of the diagnostic performance for detection of coronary artery disease (CAD) of their program (QPS) with that of the Emory Cardiac Toolbox (ECTb) for automated quantification of myocardial perfusion. <i>Journal of Nuclear Cardiology</i> , 2008 , 15, 476; author reply 476-8	2.1	3
379	Dual-energy CT of the heart for diagnosing coronary artery stenosis and myocardial ischemia-initial experience. 2008 , 18, 2414-24		187
378	Comparison of findings on stress myocardial perfusion imaging in men with versus without erectile dysfunction and without prior heart disease. 2008 , 101, 502-5		10
377	Effectiveness of anemia and chronic kidney disease as predictors for presence and severity of coronary artery disease in patients undergoing stress myocardial perfusion study. 2008 , 102, 266-71		10
376	Role of myocardial perfusion imaging in patients with end-stage renal disease undergoing coronary angiography. 2008 , 102, 1451-6		47
375	Pronfitico de la enfermedad coronaria crilica mediante gated-SPECT de perfusili miociidica. 2008 , 8, 25B-34B		
374	The diagnostic utility of the Heston index in gated SPECT to detect multi-vessel coronary artery disease. <i>Journal of Cardiology</i> , 2008 , 51, 42-9	3	2
373	Direct comparison between pharmacological stress with adenosine triphosphate disodium and exercise stress myocardial perfusion imagings. <i>Journal of Cardiology</i> , 2008 , 52, 30-8	3	4
372	High-sensitivity C-reactive protein and silent myocardial ischemia in Chinese with type 2 diabetes mellitus. 2008 , 57, 1533-8		9
371	Relationship between the QTc interval at hospital admission and the severity of the underlying ischaemia in low and intermediate risk people studied for acute chest pain. 2008 , 126, 84-91		15
370	Hypertensive response with exercise does not increase the prevalence of abnormal Tc-99m SPECT stress perfusion images. 2008 , 155, 930-7		16
369	Chronic stable angina pectoris. 2008 , 54, 671-89		8

368	Comparison of automated 4D-MSPECT and visual analysis for evaluating myocardial perfusion in coronary artery disease. 2008 , 24, 445-52	6
367	Stress Test. 2008 , 221-255	2
366	Abnormal electron beam computed tomography results: the value of repeating myocardial perfusion single-photon emission computed tomography in the ongoing assessment of coronary artery disease. 2008 , 83, 17-22	5
365	Prognostic utility of 64-slice computed tomography in patients with suspected but no documented coronary artery disease. 2009 , 30, 362-71	104
364	Quantification of regional myocardial blood flow using first-pass multidetector-row computed tomography and adenosine triphosphate in coronary artery disease. 2008 , 72, 1086-91	46
363	Incremental prognostic value of cardiac function assessed by ECG-gated myocardial perfusion SPECT for the prediction of future acute coronary syndrome. 2008 , 72, 2035-9	25
362	Myocardial perfusion imaging to assess ischemia using multislice computed tomography. 2009 , 7, 49-56	3
361	Major depression and coronary flow reserve detected by positron emission tomography. 2009 , 169, 1668-76	41
360	Normal myocardial perfusion scan portends a benign prognosis independent from the pretest probability of coronary artery disease. Sub-analysis of the J-ACCESS study. <i>Journal of Cardiology</i> , 2009 , 54, 93-100	7
359	Usefulness of abnormal heart rate recovery on exercise stress testing to predict high-risk findings on single-photon emission computed tomography myocardial perfusion imaging in men. 2009 , 103, 611-4	10
358	Comparison of dual-energy computed tomography of the heart with single photon emission computed tomography for assessment of coronary artery stenosis and of the myocardial blood supply. 2009 , 104, 318-26	150
357	Assessment of subclinical coronary atherosclerosis in asymptomatic patients with type 2 diabetes mellitus with single photon emission computed tomography and coronary computed tomography angiography. 2009 , 104, 890-6	46
356	Comparison of the value of coronary calcium detection to computed tomographic angiography and exercise testing in patients with chest pain. 2009 , 104, 1499-504	37
355	Gated myocardial perfusion SPECT asynchrony measurements in patients with left bundle branch block. 2009 , 25, 43-51	6
354	Hemodynamic variables during stress testing can predict referral to early catheterization but failed to show a prognostic impact on emerging cardiac events in patients aged 70 years and older undergoing exercise (99m)Tc-sestamibi myocardial perfusion scintigraphy. 2009 , 25, 569-79	2
353	Effect of gender on cardiovascular risk stratification with ECG gated SPECT left ventricular volume indices and ejection fraction. <i>Journal of Nuclear Cardiology</i> , 2009 , 16, 28-37	6
352	Diagnostic value of left ventricular function after adenosine triphosphate loading and at rest in the detection of multi-vessel coronary artery disease using myocardial perfusion imaging. <i>Journal of Nuclear Cardiology</i> , 2009 , 16, 20-7	17
351	Etiology and pathophysiology of new-onset heart failure: evaluation by myocardial perfusion imaging. <i>Journal of Nuclear Cardiology</i> , 2009 , 16, 82-91	41

350	Delayed heart rate recovery after adenosine stress testing with supplemental arm exercise predicts mortality. <i>Journal of Nuclear Cardiology</i> , 2009 , 16, 54-62	2.1	2
349	Threshold, incidence, and predictors of prognostically high-risk silent ischemia in asymptomatic patients without prior diagnosis of coronary artery disease. <i>Journal of Nuclear Cardiology</i> , 2009 , 16, 193	- 2d o	60
348	Detection of multivessel coronary artery disease: looking beyond the extent of perfusion abnormalities. <i>Journal of Nuclear Cardiology</i> , 2009 , 16, 4-5	2.1	3
347	Risk stratification with attenuation corrected stress Tc-99m sestamibi SPECT myocardial perfusion imaging in the absence of ECG-gating due to arrhythmias. <i>Journal of Nuclear Cardiology</i> , 2009 , 16, 533-9	2.1	26
346	Comparison of dobutamine stress echocardiography, dobutamine SPECT, and adenosine SPECT myocardial perfusion imaging in patients with end-stage renal disease. <i>Journal of Nuclear Cardiology</i> , 2009 , 16, 507-15	2.1	14
345	Stress single photon emission computed tomography for detection of coronary artery disease and risk stratification of asymptomatic patients at moderate risk. <i>Journal of Nuclear Cardiology</i> , 2009 , 16, 516-23	2.1	16
344	Prognostic validation of an algorithm to convert myocardial perfusion SPECT imaging data from a 12-segment model to a 17-segment model. <i>Journal of Nuclear Cardiology</i> , 2009 , 16, 605-13	2.1	6
343	The value of multi-slice-computed tomography coronary angiography for risk stratification. <i>Journal of Nuclear Cardiology</i> , 2009 , 16, 970-80	2.1	16
342	Interrelation of ST-segment depression during bicycle ergometry and extent of myocardial ischaemia by myocardial perfusion SPECT. 2009 , 36, 1842-50		2
341	QT variability during rest and exercise in patients with implantable cardioverter defibrillators and healthy controls. 2009 , 14, 40-9		8
340	Integrated assessment of coronary anatomy and myocardial perfusion using a retractable SPECT camera combined with 64-slice CT: initial experience. 2009 , 19, 845-56		17
339	[Agreement between observers with different experience in the interpretation of the stress myocardial perfusion electrocardiogram in SPECT]. 2009 , 28, 278-82		1
338	Myocardial perfusion, function, and dyssynchrony in patients with heart failure: baseline results from the single-photon emission computed tomography imaging ancillary study of the Heart Failure and A Controlled Trial Investigating Outcomes of Exercise TraiNing (HF-ACTION) Trial. 2009,		16
337	158, \$53-63 Prognostic value of multislice computed tomography and gated single-photon emission computed tomography in patients with suspected coronary artery disease. 2009 , 53, 623-632		272
336	Outcomes research in cardiovascular imaging: report of a workshop sponsored by the National Heart, Lung, and Blood Institute. 2009 , 2, 897-907		30
335	Agreement between observers with different experience in the interpretation of the stress myocardial perfusion electrocardiogram in SPECT. 2009 , 28, 278-282		
334	The association between erectile dysfunction and peripheral arterial disease as determined by screening ankle-brachial index testing. 2009 , 207, 440-4		44
333	Outcomes research in cardiovascular imaging: report of a workshop sponsored by the National Heart, Lung, and Blood Institute. 2009 , 3, 212-23		3

(2010-2009)

332	Prognostic value of treadmill exercise and dobutamine stress positron emission tomography. 2009 , 25, e220-4		12
331	Outcomes research in cardiovascular imaging: report of a workshop sponsored by the National Heart, Lung, and Blood Institute. 2009 , 22, 766-73		8
330	The prognostic significance of typical perfusion defects on vasodilator stress myocardial perfusion SPECT in patients with left bundle branch block or right ventricular apical pacing. 2009 , 30, 232-9		8
329	Stress-induced ST-segment deviation in relation to the presence and severity of coronary artery disease in patients with normal myocardial perfusion imaging. 2009 , 20, 41-9		
328	Sex-specific approach to gated SPECT volumetric analysis after stress and at rest to detect high-risk coronary artery disease. 2010 , 31, 800-6		
327	Rationale and design of the Randomized comparison of XiEnce V and Multilink VisioN coronary stents in the sAme multivessel patient with chronic kiDnEy disease (RENAL-DES) study. 2010 , 11, 310-7		7
326	Vasodilator stress impairs the left ventricular function obtained with gated single-photon emission computed tomography in patients with known or suspected coronary artery disease. 2010 , 74, 2666-73		3
325	Myocardial perfusion profile in a young population with and without known coronary artery disease: comparison by gender. 2010 , 33, E39-43		8
324	Comparison of diagnostic performances of three different software packages in detecting coronary artery disease. 2010 , 37, 2070-8		19
323	Stress/rest myocardial perfusion scintigraphy in patients without significant coronary artery disease. <i>Journal of Nuclear Cardiology</i> , 2010 , 17, 38-44	2.1	20
322	Direct comparison of rest and adenosine stress myocardial perfusion CT with rest and stress SPECT. Journal of Nuclear Cardiology, 2010 , 17, 27-37	2.1	83
321	Diagnostic and prognostic value of myocardial perfusion gated SPECT in orthotopic heart transplant recipients. <i>Journal of Nuclear Cardiology</i> , 2010 , 17, 197-206	2.1	30
320	The prognosis of a normal stress-only Tc-99m myocardial perfusion imaging study. <i>Journal of Nuclear Cardiology</i> , 2010 , 17, 370-7	2.1	84
319	Quantification of myocardial perfusion SPECT studies in Chinese population with Western normal databases. <i>Journal of Nuclear Cardiology</i> , 2010 , 17, 486-93	2.1	12
318	Early insights of cardiac risk and treatment response with quantitative PET monitoring of coronary-specific endothelial dysfunction and myocardial perfusion reserve. <i>Journal of Nuclear Cardiology</i> , 2010 , 17, 985-9	2.1	1
317	Does myocardial perfusion imaging provide incremental prognostic information to left ventricular ejection fraction?. <i>Current Cardiology Reports</i> , 2010 , 12, 155-61	4.2	8
316	Myocardial wall motion and thickening assessment in early gated SPECT images of acute coronary syndrome patients likely to have inferolateral perfusion defects. 2010 , 26, 881-91		2
315	The impact of chronic kidney disease as a predictor of major cardiac events in patients with no evidence of coronary artery disease. <i>Journal of Cardiology</i> , 2010 , 55, 328-36	3	16

314	Comparison of high risk stress myocardial perfusion imaging findings in men with rapid versus prolonged recovery of ST-segment depression after exercise stress testing. 2010 , 105, 1361-4	4
313	Prognostic value of a nine-minute treadmill test in patients undergoing myocardial perfusion scintigraphy. 2010 , 106, 1423-8	6
312	Prevalence, prognosis, and therapeutic implications of unrecognized left ventricular systolic dysfunction in patients with anemia and chronic kidney disease. 2010 , 16, 271-7	6
311	Prognostic value of exercise myocardial scintigraphy in patients with coronary chronic total occlusions. 2010 , 23, 139-48	24
310	Functionally unrevascularized chronic total occlusions: a step closer to indentifying the CTO patient who benefits from PCI. 2010 , 23, 149-51	
309	Coronary Artery Calcification: Pathogenesis, Imaging, and Risk Stratification. 2010 , 332-355	
308	Coronary Artery Disease Detection. 2010 , 225-243	1
307	Evaluation of plaques and stenosis. 2010 , 48, 729-44	4
306	Weighing the risks and benefits of cardiac imaging with ionizing radiation. 2010 , 3, 528-35	41
305	Renal function and risk stratification of diabetic and nondiabetic patients undergoing evaluation for coronary artery disease. 2010 , 3, 734-45	21
304	Comparison of the extent and severity of myocardial perfusion defects measured by CT coronary angiography and SPECT myocardial perfusion imaging. 2010 , 3, 1010-9	61
303	Inflammation is related to coronary flow reserve detected by positron emission tomography in asymptomatic male twins. 2011 , 57, 1271-9	54
302	Cardiac CT angiography compared with myocardial perfusion stress testing on downstream resource utilization. 2011 , 5, 101-9	22
301	CT detection of myocardial blood volume deficits: dual-energy CT compared with single-energy CT spectra. 2011 , 5, 421-9	54
300	Relationship of technetium-99m tetrofosmin-gated rest single-photon emission computed tomography myocardial perfusion imaging to death and hospitalization in heart failure patients: results from the nuclear ancillary study of the HF-ACTION trial. 2011 , 161, 1038-45	6
299	Real-world estimated effective radiation doses from commonly used cardiac testing and procedural modalities. 2011 , 27, 613-8	17
298	Vulnerable plaque features on coronary CT angiography as markers of inducible regional myocardial hypoperfusion from severe coronary artery stenoses. 2011 , 219, 588-95	65
297	Myocardial perfusion scintigraphy: techniques, interpretation, indications and reporting. 2011 , 31, 625-34	24

296	Stress testing: the case for the standard treadmill test. 2011 , 26, 363-9		12
295	Combined assessment of myocardial perfusion and function by ECG-gated myocardial perfusion single-photon emission computed tomography for the prediction of future cardiac events in patients with type 2 diabetes mellitus. 2011 , 75, 376-82		11
294	Physiological assessment of myocardial perfusion using nuclear cardiology would enhance coronary artery disease patient care: which imaging modality is best for evaluation of myocardial ischemia? (SPECT-side). 2011 , 75, 713-22; discussion 731		23
293	Cardiovascular event risk assessment using myocardial perfusion imaging with a view to wide clinical application. 2011 , 75, 2318-9		1
292	Incremental diagnostic value of regional myocardial blood flow quantification over relative perfusion imaging with generator-produced rubidium-82 PET. 2011 , 75, 2628-34		39
291	Arm exercise stress perfusion imaging predicts clinical outcome. 2011 , 111, 1546-53		5
2 90	A template-based approach to semi-quantitative SPECT myocardial perfusion imaging: Independent of normal databases. 2011 , 38, 4186-95		4
289	Myocardial perfusion imaging using adenosine-induced stress dual-energy computed tomography of the heart: comparison with cardiac magnetic resonance imaging and conventional coronary angiography. 2011 , 21, 26-35		136
288	Addition of atropine to submaximal exercise stress testing in patients evaluated for suspected ischaemia with SPECT imaging: a randomized, placebo-controlled trial. 2011 , 38, 245-51		6
287	Overall and gender-based negative predictive value of a normal gated myocardial perfusion SPECT study: a single center experience. <i>Annals of Nuclear Medicine</i> , 2011 , 25, 207-11	2.5	2
286	Clinical usefulness of combinatorial protocol with stress only myocardial perfusion SPECT, CTA and SPECT/CTA 3-dimensional fusion image. <i>Annals of Nuclear Medicine</i> , 2011 , 25, 387-95	2.5	O
285	Impact of chronic kidney disease and stress myocardial perfusion imaging as a predictor of cardiovascular events. <i>Annals of Nuclear Medicine</i> , 2011 , 25, 616-24	2.5	7
284	Correlation of chronic kidney disease, diabetes and peripheral artery disease with cardiovascular events in patients using stress myocardial perfusion imaging. <i>Annals of Nuclear Medicine</i> , 2011 , 25, 634-	4 2 5	6
283	Synergistic effect of coronary artery disease risk factors on long-term survival in patients with normal exercise SPECT studies. <i>Journal of Nuclear Cardiology</i> , 2011 , 18, 207-14; quiz 217	2.1	17
282	Non-invasive risk assessment in patients with chronic kidney disease. <i>Journal of Nuclear Cardiology</i> , 2011 , 18, 472-85	2.1	2
281	Prevalence of coronary artery calcium scores and silent myocardial ischaemia was similar in Indian Asians and European whites in a cross-sectional study of asymptomatic subjects from a U.K. population (LOLIPOP-IPC). <i>Journal of Nuclear Cardiology</i> , 2011 , 18, 435-42	2.1	5
280	Incremental prognostic value of coronary flow reserve assessed with single-photon emission computed tomography. <i>Journal of Nuclear Cardiology</i> , 2011 , 18, 541-3	2.1	1
279	Predictors of an ischemic electrocardiographic response in patients with exercise-induced myocardial ischemia. <i>Journal of Nuclear Cardiology</i> , 2011 , 18, 678-84	2.1	2

278	Small average differences in attenuation corrected images between men and women in myocardial perfusion scintigraphy: a novel normal stress database. 2011 , 11, 18	4
277	One-year prognosis of patients with normal myocardial perfusion imaging using technitium-99m sestamibi in suspected coronary artery disease: a single-center experience of 1,047 patients. 2011 , 26, 309-14	2
276	Role of Stress Myocardial Scintigraphy in the Evaluation of Incompletely Revascularized Post-PCI Patients. 2011 , 2011, 180936	1
275	Assessment of cardiac ischaemia and viability: role of cardiovascular magnetic resonance. 2011 , 32, 799-809	57
274	Diabetes reduces left ventricular ejection fractionirrespective of presence and extent of coronary artery disease. 2011 , 165, 945-51	18
273	Prognostic value of adenosine stress myocardial perfusion by cardiac magnetic resonance imaging in patients with known or suspected coronary artery disease. 2011 , 104, 425-32	11
272	Diagnostic performance of combined noninvasive anatomic and functional assessment with dual-source CT and adenosine-induced stress dual-energy CT for detection of significant coronary stenosis. 2012 , 198, 512-20	95
271	Adenosine-stress dynamic myocardial perfusion imaging with second-generation dual-source CT: comparison with conventional catheter coronary angiography and SPECT nuclear myocardial perfusion imaging. 2012 , 198, 521-9	110
270	Toward a practical template-based approach to semiquantitative SPECT myocardial perfusion imaging. 2012 , 39, 1374-85	0
269	Prognostic implications of the difference between left ventricular ejection fractions after stress and at rest in addition to the quantification of myocardial perfusion abnormalities obtained with gated SPECT. 2012 , 37, 748-54	7
268	CT Evaluation of the Myocardial Blood Supply: Dual-Source Dual-Energy CT. 2012 , 79-102	
267	Postischemic myocardial stunning is superior to transient ischemic dilation for detecting multivessel coronary artery disease. 2012 , 76, 430-8	11
266	2012 ACCF/AHA/ACP/AATS/PCNA/SCAI/STS Guideline for the diagnosis and management of patients with stable ischemic heart disease: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines, and the American	1138
265	College of Physicians, American Association for Thoracic Surgery, Preventive Cardiovascular Nurses Prognostic value of coronary artery calcium score and coronary CT angiography in patients with intermediate risk of coronary artery disease. 2012, 28, 1547-56	32
264	Risk stratification of cardiovascular events in patients at all stages of chronic kidney disease using myocardial perfusion SPECT. <i>Journal of Cardiology</i> , 2012 , 60, 377-82	27
263	Downstream procedures and outcomes after stress testing for chest pain without known coronary artery disease in the United States. 2012 , 163, 454-61	29
262	Nuclear medicine technologists are able to accurately determine when a myocardial perfusion rest study is necessary. 2012 , 12, 97	2
261	Polymeric Imaging Agents. 2012 , 529-545	

260	Normal stress databases in myocardial perfusion scintigraphyhow many subjects do you need?. 2012 , 32, 455-62		1
259	Patient management after noninvasive cardiac imaging results from SPARC (Study of myocardial perfusion and coronary anatomy imaging roles in coronary artery disease). 2012 , 59, 462-74		143
258	Journey in evolution of nuclear cardiology: will there be another quantum leap with the F-18-labeled myocardial perfusion tracers?. 2012 , 5, 1269-84		47
257	Adenosine-stress dynamic real-time myocardial perfusion CT and adenosine-stress first-pass dual-energy myocardial perfusion CT for the assessment of acute chest pain: initial results. 2012 , 81, 3703-10		106
256	Safety and feasibility of regadenoson use for suboptimal heart rate response during symptom-limited standard Bruce exercise stress test. <i>Journal of Nuclear Cardiology</i> , 2012 , 19, 970-8	2.1	21
255	Long-term prognostic value of exercise technetium-99m tetrofosmin myocardial perfusion single-photon emission computed tomography. <i>Journal of Nuclear Cardiology</i> , 2012 , 19, 907-13	2.1	18
254	Prognostic value of myocardial perfusion scintigraphy in elderly patients with hypertension: a 10-year follow-up analysis. 2012 , 39, 1570-80		3
253	Referring physicians underestimate the extent of abnormalities in final reports from myocardial perfusion imaging. 2012 , 2, 27		5
252	Safety and efficacy of exercise testing with atropine in patients with recent uncomplicated ST elevation acute myocardial infarction. 2012 , 7, 262-268		
251	Effect of vascular endothelial growth factor gene transfer on infarct size, left ventricular function and myocardial perfusion in sheep after 2 months of coronary artery occlusion. 2012 , 14, 279-87		11
250	Body mass index and rest myocardial perfusion defect predicts cardiac death in patients with chronic heart failure. 2012 , 28 Suppl 1, 77-85		5
249	Does location matter? Prognostic value of single-photon emission computed tomography myocardial perfusion imaging by vascular territory. <i>Journal of Nuclear Cardiology</i> , 2012 , 19, 458-64	2.1	3
248	Stress CT perfusion: coupling coronary anatomy with physiology. <i>Journal of Nuclear Cardiology</i> , 2012 , 19, 588-600	2.1	21
247	Use of high-risk features from exercise treadmill testing to identify obstructive left main disease with normal myocardial perfusion imaging. <i>Journal of Nuclear Cardiology</i> , 2012 , 19, 814-7	2.1	
246	Effects of type 2 diabetes mellitus on coronary microvascular function and myocardial perfusion in patients without obstructive coronary artery disease. 2012 , 39, 1199-206		44
245	Utility of cardiac computed tomography angiography to exclude clinically significant obstructive coronary artery disease in patients after myocardial perfusion imaging. 2012 , 109, 165-8		19
244	The complementary roles of radionuclide myocardial perfusion imaging and cardiac computed tomography. 2012 , 47, 228-39		3
243	Selecting the best noninvasive imaging test to guide treatment after an inconclusive exercise test. 2012 , 14, 8-23		4

242	Severity of inducible myocardial ischemia predicts incident acute coronary syndromes in asymptomatic individuals with a family history of premature coronary artery disease. <i>Journal of Nuclear Cardiology</i> , 2012 , 19, 28-36	11
241	How to evaluate physical fitness without a stress test?. 2012 , 28, 199-209	
240	When is reacquisition necessary due to high extra-cardiac uptake in myocardial perfusion scintigraphy?. 2013 , 3, 20	4
239	Population-based study of the use of cardiac stress imaging and referral for coronary angiography and repeated revascularization after coronary artery bypass graft surgery. 2013 , 88, 345-53	4
238	Adding attenuation corrected images in myocardial perfusion imaging reduces the need for a rest study. 2013 , 13, 14	8
237	Validation of videodensitometric myocardial perfusion assessment. 2013 , 8, 600-607	
236	A hypothetical protocol for the provisional use of perfusion imaging with exercise stress testing. Journal of Nuclear Cardiology, 2013 , 20, 739-47	10
235	Comparison of attenuation, dual-energy-window, and model-based scatter correction of low-count SPECT to 82Rb PET/CT quantified myocardial perfusion scores. <i>Journal of Nuclear Cardiology</i> , 2013 , 21, 20, 785-96	5
234	The prognosis for prognosis remains excellent. <i>Journal of Nuclear Cardiology</i> , 2013 , 20, 501-3	1
233	The prognostic value of cardiac SPECT performed at the primary care physician's office. <i>Journal of Nuclear Cardiology</i> , 2013 , 20, 519-28	27
232	Microalbuminuria predicts silent myocardial ischaemia in type 2 diabetes patients. 2013 , 40, 548-57	9
231	Prognosis of patients without perfusion defects with and without rest study in myocardial perfusion scintigraphy. 2013 , 3, 58	9
230	Downstream clinical implications of abnormal myocardial perfusion single-photon emission computed tomography based on appropriate use criteria. <i>Journal of Nuclear Cardiology</i> , 2013 , 20, 1041-8 ^{2.1}	14
229	Prediction of 8-year cardiovascular outcomes in patients with systemic arterial hypertension: value of stress (99m)Tc-tetrofosmin myocardial perfusion imaging in a high-risk cohort. <i>Journal of Nuclear 2.1 Cardiology</i> , 2013 , 20, 1030-40	3
228	Predictors of cardiovascular events in hemodialysis patients after stress myocardial perfusion imaging. 2013 , 17, 568-75	1
227	The predictive value of chronic kidney disease for assessing cardiovascular events under consideration of pretest probability for coronary artery disease in patients who underwent stress myocardial perfusion imaging. 2013 , 29, 513-9	2
226	Comparison of the applicability of the 2006 and the 2010 cardiac CT angiography appropriate use criteria. 2013 , 10, 258-67	4
225	Repeated sauna therapy improves myocardial perfusion in patients with chronically occluded coronary artery-related ischemia. 2013 , 167, 237-43	37

224	The role of noninvasive imaging in coronary artery disease detection, prognosis, and clinical decision making. 2013 , 29, 285-96	28
223	Myocardial perfusion imaging with 13N-ammonia PET is a strong predictor for outcome. 2013 , 167, 1023-6	29
222	Prognosis of patients with mild-moderate coronary artery stenosis detected by coronary computed tomography angiography. 2013 , 168, 1195-200	14
221	Relationship between silent myocardial ischemia and coronary artery disease risk factors. <i>Journal of Nuclear Cardiology</i> , 2013 , 20, 731-8	16
220	Value of reserve pulse pressure in improving the risk stratification of patients with normal myocardial perfusion imaging. 2013 , 34, 2074-81a	4
219	Vitamin D status and coronary flow reserve measured by positron emission tomography: a co-twin control study. 2013 , 98, 389-97	12
218	Detection of ischaemic myocardial lesions with coronary CT angiography and adenosine-stress dynamic perfusion imaging using a 128-slice dual-source CT: diagnostic performance in comparison with cardiac MRI. 2013 , 86, 20130481	16
217	Stress cardiac magnetic resonance imaging provides effective cardiac risk reclassification in patients with known or suspected stable coronary artery disease. <i>Circulation</i> , 2013 , 128, 605-14	50
216	Long-term prognostic value of inducible and resting perfusion defects detected by single-photon emission computed tomography in the era of wide availability of coronary revascularization. 2013 , 33, 218-23	6
215	Gender differences in myocardial perfusion defect in asymptomatic postmenopausal women and men with and without diabetes mellitus. 2013 , 22, 439-44	8
214	Detection of post-exercise stunning by early gated SPECT myocardial perfusion imaging: results from the IAEA multi-center study. <i>Journal of Nuclear Cardiology</i> , 2014 , 21, 1168-76	21
213	Prognostic value of regadenoson myocardial single-photon emission computed tomography in patients with different degrees of renal dysfunction. 2014 , 15, 933-40	19
212	Endothelial Dysfunction is Associated with Occult Coronary Artery Disease Detected by Positron Emission Tomography. 2014 , 4, 28-32	4
211	Computer-aided diagnosis system outperforms scoring analysis in myocardial perfusion imaging. <i>Journal of Nuclear Cardiology</i> , 2014 , 21, 416-23	14
210	Prognostic role of stress/rest myocardial perfusion scintigraphy in patients with cardiac syndrome x. 2014 , 173, 467-71	6
209	Assessing clinical impact of myocardial perfusion studies: ischemia or other prognostic indicators?. Current Cardiology Reports, 2014 , 16, 465 4.2	1
208	Area of ischemia assessed by physicians and software packages from myocardial perfusion scintigrams. 2014 , 14, 5	6
207	The role of nuclear cardiology in the diagnosis and risk stratification of women with ischemic heart disease. 2014 , 44, 423-38	3

206	Resultados de SPECT miocfidico y eventos cardiovasculares en dos servicios de Medicina nuclear de Bogotfidurante 2011 a 2012. 2014 , 21, 86-94		1
205	Noninvasive stress testing for coronary artery disease. 2014 , 32, 387-404		7
204	Clinical decision making with myocardial perfusion imaging in patients with known or suspected coronary artery disease. 2014 , 44, 320-9		38
203	Direct comparison of stress- and rest-dual-energy computed tomography for detection of myocardial perfusion defect. 2014 , 30 Suppl 1, 41-53		41
202	Novel insights into the pathophysiology of different forms of stress testing. 2014 , 47, 338-43		6
201	New insights from major prospective cohort studies with cardiac nuclear imaging. <i>Current Cardiology Reports</i> , 2014 , 16, 482	4.2	1
200	Risk stratification using line source attenuation correction with rest/stress Tc-99m sestamibi SPECT myocardial perfusion imaging. <i>Journal of Nuclear Cardiology</i> , 2014 , 21, 118-26	2.1	14
199	B-type natriuretic peptide and clinical judgment in the detection of exercise-induced myocardial ischemia. 2014 , 127, 427-35		15
198	Comment on âDirect comparison of cardiac magnetic resonance and multidetector computed tomography stress-rest perfusion imaging for detection of coronary artery diseaseâ[]2014, 33, 125-126		
197	Comentiio a «Comparati direta entre a ressoniicia magntica cardâca e a tomografia computorizada multidetetores de perfusti na deteti de doenii coroniia». 2014 , 33, 125-126		
196	Prediction of cardiovascular events in pre-dialysis chronic kidney disease patients with normal SPECT myocardial perfusion imaging. <i>Journal of Cardiology</i> , 2014 , 63, 154-8	3	7
195	Usefulness of stress myocardial perfusion imaging and baseline clinical factors for predicting cardiovascular events in patients with prior coronary artery disease. 2014 , 78, 1676-83		7
194	Perfusion vector-a new method to quantify myocardial perfusion scintigraphy images: a simulation study with validation in patients. 2015 , 5, 121		1
193	The Significance of ST Depression in a Postmenopausal Woman on Estrogen Therapy during Regadenoson Myocardial SPECT Imaging. 2015 , 2015, 653760		1
192	Myocardial Perfusion Imaging Utilizing Single Photon Emission Computed Tomography Techniques. 2015 , 253-262		
191	A randomized, multicenter, multivendor study of myocardial perfusion imaging with regadenoson CT perfusion vs single photon emission CT. 2015 , 9, 103-12.e1-2		56
190	Evaluation of inter-departmental variability of ejection fraction and cardiac volumes in myocardial perfusion scintigraphy using simulated data. 2015 , 2, 2		5
189	Prediction of 14-year outcomes in patients with a limited exercise capacity: Utility of dobutamine myocardial perfusion imaging in a high-risk population. <i>Journal of Nuclear Cardiology</i> , 2015 , 22, 888-900	2.1	О

188	Incremental value of a single high-sensitivity cardiac troponin I measurement to rule out myocardial ischemia. 2015 , 128, 638-46	25
187	Revascularization vs. Medical Therapy in Stable Ischemic Heart Disease. 2015 , 58, 299-305	6
186	Eleven-year prognostic value of dobutamine stress (99m)Tc-sestamibi myocardial perfusion imaging in patients with limited exercise capacity. 2015 , 115, 884-9	10
185	Radionuclide myocardial perfusion imaging for the evaluation of patients with known or suspected coronary artery disease in the era of multimodality cardiovascular imaging. 2015 , 57, 644-53	19
184	Added prognostic value of ischaemic threshold in radionuclide myocardial perfusion imaging: a common-sense integration of exercise tolerance and ischaemia severity. 2015 , 42, 750-60	4
183	Effective risk stratification of patients on the basis of myocardial perfusion SPECT is dependent on appropriate patient selection. <i>Current Cardiology Reports</i> , 2015 , 17, 549	6
182	Scatter correction improves concordance in SPECT MPI with a dedicated cardiac SPECT solid-state camera. <i>Journal of Nuclear Cardiology</i> , 2015 , 22, 334-43	19
181	An Increasing Population with Metabolic Syndrome and/or Diabetes Mellitus in the Middle Eastâls There an Added Value of Coronary Calcium Scoring to Myocardial Perfusion Imaging?. 2015 , 8, 1	
180	Serious and potentially life threatening complications of cardiac stress testing: Physiological mechanisms and management strategies. <i>Journal of Nuclear Cardiology</i> , 2015 , 22, 1198-213; quiz 1195-7 ^{2.1}	21
179	Use of Technetium-99m-Labeled Lipophilic Complex of Methoxy-Isobutyl-Isonitrile and Iopromide Radiopaque Substance to Assess the Severity of Coronary Atherosclerosis in Patients with Coronary Heart Disease at Mild Disorders of Myocardial Perfusion. 2015 , 1084, 527-531	
178	Significance of worsening renal function and nuclear cardiology for predicting cardiac death in patients with known or suspected coronary artery disease. <i>Journal of Cardiology</i> , 2015 , 66, 423-9	4
177	Prognostic Contribution of Exercise Capacity, Heart Rate Recovery, Chronotropic Incompetence, and Myocardial Perfusion Single-Photon Emission Computerized Tomography in the Prediction of Cardiac Death and All-Cause Mortality. 2015 , 116, 1678-84	19
176	Application of the newly developed Japanese adenosine normal database for adenosine stress myocardial scintigraphy. <i>Annals of Nuclear Medicine</i> , 2015 , 29, 730-9	
175	Comparison of the prognostic value of regadenoson and adenosine myocardial perfusion imaging. <i>Journal of Nuclear Cardiology</i> , 2015 , 22, 600-7	18
174	Diagnostic and Prognostic Role of Myocardial Perfusion Scintigraphy in Kidney Transplant Candidates: Narrative Review. 2016 , 11, heartint.500023	
173	Revascularization for Silent Myocardial Ischemia. 2016 , 111-124	
172	Adult Gamma Camera Myocardial Perfusion Imaging: Diagnostic Reference Levels and Achievable Administered Activities Derived From ACR Accreditation Data. 2016 , 13, 688-95	9
171	Usefulness of the novel risk estimation software, Heart Risk View, for the prediction of cardiac events in patients with normal myocardial perfusion SPECT. <i>Annals of Nuclear Medicine</i> , 2016 , 30, 716-721.5	2

170	Differential risk reclassification improvement by exercise testing and myocardial perfusion imaging in patients with suspected and known coronary artery disease. <i>Journal of Nuclear Cardiology</i> , 2016 , 23, 366-78	2.1	6
169	Clinical Feasibility of Simultaneous Acquisition Rest (99m)Tc/Stress (201)Tl Dual-Isotope Myocardial Perfusion Single-Photon Emission Computed Tomography With Semiconductor Camera. 2016 , 80, 689-9	95	21
168	The Sphinx's riddle: cardiovascular involvement in autoimmune rheumatic disease. 2016 , 16, 204		5
167	Exercise and Bayes' Theorem: Some things never go out of style. <i>Journal of Nuclear Cardiology</i> , 2016 , 23, 379-83	2.1	1
166	Prediction Models for Cardiac Risk Classification with Nuclear Cardiology Techniques. 2016 , 9, 1		
165	Temporal trends of single-photon emission computed tomography myocardial perfusion imaging in patients without prior coronary artery disease: A 22-year experience at a tertiary academic medical center. 2016 , 176, 127-33		20
164	The "rise of the rule out": The SPECTer of cardiac stress imaging. 2016 , 176, 122-4		
163	Insulin resistance, endothelial function, angiogenic factors and clinical outcome in non-diabetic patients with chest pain without myocardial perfusion defects. 2016 , 15, 36		16
162	Different prognosis according to different clinical, electrocardiographic and scintigraphic ischemia criteria. 2016 , 219, 240-6		3
161	Dobutamine stress myocardial perfusion imaging: 8-year outcomes in patients with diabetes mellitus. 2016 , 17, 871-6		4
160	Diagnostic Strategies for the Evaluation of Chest Pain: Clinical Implications From SCOT-HEART and PROMISE. 2016 , 67, 843-52		41
159	Optimal non-invasive imaging test selection for the diagnosis of ischaemic heart disease. <i>Heart</i> , 2016 , 102, 555-64	5.1	9
158	Cardiovascular outcomes after pharmacologic stress myocardial perfusion imaging. 2016 , 174, 138-46		4
157	Direct comparison of cardiac troponin I and cardiac troponin T in the detection of exercise-induced myocardial ischemia. 2016 , 49, 421-432		17
156	Software-based on-site estimation of fractional flow reserve using standard coronary CT angiography data. 2016 , 57, 1186-92		34
155	Noninvasive Stress Testing for Coronary Artery Disease. 2016 , 12, 65-82		7
154	Prognostic evaluation in obese patients using a dedicated multipinhole cadmium-zinc telluride SPECT camera. 2016 , 32, 355-361		12
153	Clinical value of high duke treadmill score with myocardial perfusion SPECT. <i>Journal of Nuclear Cardiology</i> , 2016 , 23, 1301-1303	2.1	4

152	Stress testing in patients with chronic kidney disease: The need for ancillary markers for effective risk stratification and prognosis. <i>Journal of Nuclear Cardiology</i> , 2016 , 23, 570-4	2.1	15
151	Prognostic value of the cadmium-zinc-telluride camera: A comparison with a conventional (Anger) camera. <i>Journal of Nuclear Cardiology</i> , 2016 , 23, 1280-1287	2.1	12
150	Stress protocol and accuracy of myocardial perfusion imaging: Is it better to start from the end?. <i>Journal of Nuclear Cardiology</i> , 2016 , 23, 1123-1127	2.1	2
149	Quantitative iodine-123-metaiodobenzylguanidine (MIBG) SPECT imaging in heart failure with left ventricular systolic dysfunction: Development and validation of automated procedures in conjunction with technetium-99m tetrofosmin myocardial perfusion SPECT. <i>Journal of Nuclear</i>	2.1	19
148	Diagnostic and Prognostic Value of Lead aVR During Exercise Testing in Patients Suspected of Having Myocardial Ischemia. 2017 , 119, 959-966		8
147	Global quantification of left ventricular myocardial perfusion at dynamic CT imaging: Prognostic value. 2017 , 11, 16-24		17
146	Stable CAD and Approach to Chronic Chest Pain. 2017 , 65-91		
145	Stress Testing, Nuclear Imaging, Coronary CT Angiography. 2017 , 649-664		
144	Usefulness for Predicting Cardiac Events After Orthotopic Liver Transplantation of Myocardial Perfusion Imaging and Dobutamine Stress Echocardiography Preoperatively. 2017 , 119, 1008-1011		5
143	The diagnostic accuracy of myocardial perfusion scintigraphy in athletes with abnormal exercise test results. 2017 , 24, 1000-1007		3
142	Prognostic Value of Stress Dynamic Myocardial Perfusion CT in a Multicenter Population With Known or Suspected Coronary Artery Disease. 2017 , 208, 761-769		19
141	Outcomes-Based CV Imaging Research Endpoints and Trial Design: From Pixels to Patient Satisfaction. 2017 , 10, 253-263		2
140	Identification of Patients With Stable Chest Pain Deriving Minimal Value From Noninvasive Testing: The PROMISE Minimal-Risk Tool, A Secondary Analysis of a Randomized Clinical Trial. 2017 , 2, 400-408		44
139	Prognostic value of normal positron emission tomography myocardial perfusion imaging in patients with known or suspected coronary artery disease: a meta-analysis. 2017 , 90, 20160702		4
138	CT myocardial perfusion imaging: current status and future perspectives. 2017 , 33, 1009-1020		13
137	New solid state cadmium-zinc-telluride technology for cardiac single photon emission computed tomographic myocardial perfusion imaging. 2017 , 14, 213-222		8
136	Detection of Myocardial Infarction Using Delayed Enhancement Dual-Energy CT in Stable Patients. 2017 , 209, 1023-1032		7
135	The elusive role of myocardial perfusion imaging in stable ischemic heart disease: Is ISCHEMIA the answer?. <i>Journal of Nuclear Cardiology</i> , 2017 , 24, 1610-1618	2.1	2

Risk stratification of coronary artery disease using radionuclides. Current status of clinical practice. **2017**, 36, 377-387

133	Risk stratification of coronary artery disease using radionuclides. Current status of clinical practice. 2017 , 36, 377-387		
132	What do we know? What do we need to know?. Journal of Nuclear Cardiology, 2017, 24, 252-254	2.1	7
131	Cardiovascular risk stratification after renal transplant: Is SPECT-MPI the answer?. <i>Journal of Nuclear Cardiology</i> , 2017 , 24, 304-307	2.1	1
130	Revascularization and outcomes in Veterans with moderate to severe ischemia on myocardial perfusion imaging. 2017 , 4, 12		1
129	Has anyone been listening? Post-SPECT MPI referral rates to catheterization. <i>Journal of Nuclear Cardiology</i> , 2017 , 24, 1662-1665	2.1	1
128	Functional compared to anatomical imaging in the initial evaluation of patients with suspected coronary artery disease: An international, multi-center, randomized controlled trial (IAEA-SPECT/CTA study). <i>Journal of Nuclear Cardiology</i> , 2017 , 24, 507-517	2.1	21
127	Spect image reconstruction under imaging time constraints. 2017 ,		
126	Impact of Follow-Up Ischemia on Myocardial Perfusion Single-Photon Emission Computed Tomography in Patients with Coronary Artery Disease. 2017 , 58, 934-943		2
125	Cardiac stress imaging for the prediction of very long-term outcomes: Dobutamine stress echocardiography or dobutamine Tc-sestamibi SPECT?. <i>Journal of Nuclear Cardiology</i> , 2018 , 25, 471-479	2.1	O
124	Prediction of 14-year cardiovascular outcomes by dobutamine stress Tc-tetrofosmin myocardial perfusion SPECT in elderly patients unable to perform exercise testing. <i>Journal of Nuclear Cardiology</i> , 2018 , 25, 63-71	2.1	3
123	Fully automated analysis of attenuation-corrected SPECT for the long-term prediction of acute myocardial infarction. <i>Journal of Nuclear Cardiology</i> , 2018 , 25, 1353-1360	2.1	11
122	Updating Algorithms for Predicting Pre-Test Likelihood of Coronary´Artery´Disease: A Cure for Inappropriate Testing?. 2018 , 11, 447-449		1
121	Visualization of the improvement of myocardial perfusion after coronary intervention using motorized fractional flow reserve pullback curve. 2018 , 33, 99-108		8
120	Prognostic Value of Combined Clinical and Myocardial Perfusion Imaging Data Using Machine Learning. 2018 , 11, 1000-1009		99
119	Prognostic Impact of Extent, Severity, and Heterogeneity of Abnormalities on F-FDG PET Scans for Suspected Cardiac Sarcoidosis. 2018 , 11, 336-345		51
118	Impact of initial myocardial perfusion imaging versus invasive coronary angiography on outcomes in coronary artery disease: a nationwide cohort study. 2018 , 45, 567-574		7
117	Deciding wisely: A case for an effective use of myocardial perfusion imaging. <i>Journal of Nuclear Cardiology</i> , 2018 , 25, 53-61	2.1	О

116	Relationship between left ventricular dyssynchrony and scar burden in the genesis of ventricular tachyarrhythmia. <i>Journal of Nuclear Cardiology</i> , 2018 , 25, 555-569	2.1	5
115	Putting It All Together: Which Test for Which Patient?. 2018 , 204-225		1
114	. 2018,		O
113	Coronary pressure-derived parameters. 2018 , 26, 375-376		
112	Diagnostic Performance of the Simultaneous Acquisition Rest Tc-Tetrofosmin/Stress Tl Dual-Isotope Protocol With a Semiconductor Camera - Comparison With the Rest-Stress Tc-Tetrofosmin Protocol. 2018 , 82, 2837-2844		2
111	Single Photon Emission Computed Tomography (SPECT) Myocardial Perfusion Imaging Guidelines: Instrumentation, Acquisition, Processing, and Interpretation. <i>Journal of Nuclear Cardiology</i> , 2018 , 25, 1784-1846	2.1	126
110	Receipt of cardiac screening does not influence 1-year post-cerebrovascular event mortality. 2018 , 8, 192-200		2
109	Using time series analysis to predict affecting factors of thallium myocardial perfusion scan service usage frequency. 2018 ,		2
108	International Study of Comparative Health Effectiveness with Medical and Invasive Approaches-Chronic Kidney Disease (ISCHEMIA-CKD): Rationale and design. 2018 , 205, 42-52		27
107	Differences in attenuation pattern in myocardial SPECT between CZT and conventional gamma cameras. <i>Journal of Nuclear Cardiology</i> , 2019 , 26, 1984-1991	2.1	10
106	Reducing radiation dose from myocardial perfusion imaging in subjects with complex congenital heart disease. <i>Journal of Nuclear Cardiology</i> , 2021 , 28, 1395-1408	2.1	6
105	Risk Stratification of Cardiovascular Events in Very Elderly Patients with Known or Suspected Coronary Artery Disease Who Had Normal Single-photon Emission Computed Tomographic Myocardial Perfusion Imaging Findings. 2019 , 58, 3351-3359		
104	Algorithmic Versus Expert Human Interpretation of Instantaneous Wave-Free Ratio Coronary Pressure-Wire Pull Back Data. 2019 , 12, 1315-1324		13
103	The effect of cardiac shock wave therapy on myocardial function and perfusion in the randomized, triple-blind, sham-procedure controlled study. 2019 , 17, 13		5
102	Controversies in Diagnostic Imaging of Patients With Suspected Stable and Acute Chest Pain Syndromes. 2019 , 12, 1254-1278		3
101	Effects of silent myocardial ischemia on functional fitness and physical independence in 60â🛮 9-year-old adults. 2019 , 1, 44-48		
100	Quantification of myocardial blood flow by CZT-SPECT with motion correction and comparison with O-water PET. <i>Journal of Nuclear Cardiology</i> , 2021 , 28, 1477-1486	2.1	14
99	Diagnosis and Prognosis of Coronary Artery Disease with SPECT and PET. <i>Current Cardiology Reports</i> , 2019 , 21, 57	4.2	14

98	Choosing the Appropriate Stress Test for Myocardial Perfusion Imaging. 2019, 12, 1		1
97	A Novel Technology of Multi-pinhole SPECT for Human Myocardial Perfusion Imaging. 2019,		O
96	Fragmented QRS complex is an independent predictor of plaque burden in patients at intermediate risk of coronary artery disease. 2019 , 71, 394-399		2
95	Myocardial perfusion imaging and coronary calcium score: A marriage made in heaven. <i>Journal of Nuclear Cardiology</i> , 2021 , 28, 2097-2099	2.1	2
94	Coronary calcium score influences referral for invasive coronary angiography after normal myocardial perfusion SPECT. <i>Journal of Nuclear Cardiology</i> , 2019 , 26, 602-612	2.1	2
93	How variable are the volumetric measurements from gated perfusion SPECT when a one-day stress-rest protocol is used?. <i>Journal of Nuclear Cardiology</i> , 2019 , 26, 1709-1716	2.1	O
92	Evaluation of the effect of reducing administered activity on assessment of function in cardiac gated SPECT. <i>Journal of Nuclear Cardiology</i> , 2020 , 27, 562-572	2.1	4
91	CRAX: A simple cardiovascular risk assessment tool to predict risk of acute myocardial infarction or death. <i>Journal of Nuclear Cardiology</i> , 2020 , 27, 2365-2374	2.1	4
90	5-Year Prognostic Value of Quantitative Versus Visual MPI in Subtle Perfusion Defects: Results From REFINE SPECT. 2020 , 13, 774-785		29
89	Association Between Change in Circulating Progenitor Cells During Exercise Stress and Risk of Adverse Cardiovascular Events in Patients With Coronary Artery Disease. 2020 , 5, 147-155		6
88	Old unsolved problems: when and how to treat silent ischaemia. 2020 , 22, L82-L85		О
87	ISCHEMIA questions and MITNEC answers: Defining and standardizing clinical ischemic jeopardy with SPECT myocardial perfusion imaging. <i>Journal of Nuclear Cardiology</i> , 2020 , 1	2.1	
86	Long-Term Prognostic Value of Coronary Atherosclerotic Plaque Characteristics Assessed by Computerized Tomographic Angiography. 2021 , 72, 252-259		1
85	Quantitation in Nuclear Cardiac Imaging. 2021 , 501-530		
84	Drawing the line between a normal and mildly abnormal nuclear cardiology scan. <i>Journal of Nuclear Cardiology</i> , 2021 , 1	2.1	0
83	Prognostic Performance of Myocardial Perfusion and Function. 2021 , 325-368		
82	Exercise and Pharmacologic Stress Testing. 2021 , 517-544		
81	Approach to stable angina in patients with advanced chronic kidney disease. 2021 , 30, 339-345		1

80	Stress Myocardial Perfusion Imaging Interpretation From the Viewpoint of Fractional Flow Reserve. 2021 , 85, 2043-2049		1
79	Visually estimated coronary artery calcium score improves SPECT-MPI risk stratification. 2021 , 35, 100827		4
78	Experimental examination of radiation doses from cardiac and liver CT perfusion in a phantom study as a function of organ, age and sex. 2021 , 41,		O
77	Stable Ischemic Heart Disease. 2021 , 109-140		
76	Risk Stratification and Patient Management. 2013 , 247-288		1
75	Perfusion Measurements of the Myocardium. 2015 , 1279-1354		1
74	Diagnostic strategies for women with suspected coronary artery disease. 2001 , 15, 39-53		8
73	Prognostic value of dobutamine stress myocardial perfusion echocardiography in patients with known or suspected coronary artery disease and normal left ventricular function. <i>PLoS ONE</i> , 2017 , 12, e0172280	7	4
72	"Save the Last Dance" for Cardiovascular Magnetic Resonance. <i>European Cardiology Review</i> , 2018 , 13, 95-97	9	1
71	Prognostic Assessment of Diabetics Using Myocardial Perfusion Imaging: Diabetes Mellitus is Still a Coronary Artery Disease Equivalent. <i>Open Cardiovascular Medicine Journal</i> , 2017 , 11, 76-83	7	2
70	Appropriate Use Criteria for PET Myocardial Perfusion Imaging. <i>Journal of Nuclear Medicine</i> , 2020 , 61, 1221-1265	9	16
69	Update of the Brazilian Guideline on Nuclear Cardiology - 2020. <i>Arquivos Brasileiros De Cardiologia</i> , 2020 , 114, 325-429	2	4
68	Long-term prognosis of end-stage renal disease patients with normal myocardial perfusion as determined by single photon emission computed tomography. <i>Korean Journal of Internal Medicine</i> , 2. 2018 , 33, 148-156	5	3
67	Assessment of prognostic value of semiquantitative parameters on gated single photon emission computed tomography myocardial perfusion scintigraphy in a large middle eastern population. O. Indian Journal of Nuclear Medicine, 2015, 30, 233-8	4	4
66	Effectiveness of (99m)Tc-tetrofosmin for assessment of heart functions in micropigs. <i>Journal of Veterinary Science</i> , 2007 , 8, 223-7	6	1
65	Myocardial Segmentation of Area at Risk Based on Coronary Computed Tomography Angiography and Voronoi Diagram in Comparison with Magnetic Resonance Perfusion Imaging. <i>Open Journal of Radiology</i> , 2017 , 07, 9-22	2	2
64	Transcatheter treatment for refractory angina with the Coronary Sinus Reducer. <i>EuroIntervention</i> , 2014 , 9, 1158-64	1	36
63	Myocardial ischemia is a key factor in the management of stable coronary artery disease. <i>World Journal of Cardiology</i> , 2014 , 6, 130-9	1	17

62	Myocardium in Jeopardy. Developments in Cardiovascular Medicine, 2001, 119-144			
61	Interpreting Noninvasive Cardiac Tests. 2001 , 25-36			
60	Effect of power Doppler and digital subtraction techniques on the comparison of myocardial contrast echocardiography with SPECT. <i>Heart</i> , 2001 , 85, 549-555	5.1		
59	Gated Myocardial Perfusion Spect for Diagnosis and Prognosis of Patients with Coronary Artery Disease. <i>Developments in Cardiovascular Medicine</i> , 2002 , 137-155			
58	Novel Internet-based tool for correcting apparent sensitivity and specificity of diagnostic tests to adjust for referral (verification) bias. <i>Radiographics</i> , 2002 , 22, e4	5.4	1	
57	Risk Stratification and Patient Management. 2003 , 97-113			
56	Stress Modalities to Evaluate Myocardial Perfusion. 2004 , 84-120			
55	Should All Women With Suspected Coronary Disease Undergo Stress Cardiac Imaging?. 2004 , 161-172			
54	Myocardial Perfusion Imaging for Cardiac Risk Stratification. 2004 , 253-278			
53	Applications of Myocardial Perfusion Imaging in Special Populations. 2004 , 331-344			
52	Nuclear Cardiology for Imaging the Effects of Therapy. 2004 , 279-308			
51	Medicina basada en la evidencia como guâ para el diagnEtico por imagen del corazE. 2005 , 186-196			
50	Diagnosis of Coronary Artery Disease. 2006 , 155-187			
49	Prognostic Assessment by Noninvasive Imaging. Part a. Clinical Decision-making in Patients with Suspected or Known Coronary Artery Disease. 2006 , 189-208			
48	Myocardial Perfusion Imaging Utilizing Single Photon Emission Computed Tomography Techniques. 2007 , 841-853			
47	Cardiovascular Imaging in Racial/Ethnic Populations: Implications for the Adequate Application of Cardiovascular Imaging Techniques Guided by Racial/Ethnic Risk Factor Variations. 2009 , 229-245			
46	Comparative Use of Radionuclide Stress Testing, Coronary Artery Calcium Scanning, and Noninvasive Coronary Angiography for Diagnostic and Prognostic Cardiac Assessment. 2010 , 233-254			

Quantitative Cardiac SPECT Imaging. **2010**, 311-331

Prognostic Implications of MPI Stress SPECT. 2010, 267-286 44 Nuclear Cardiology and Positron Emission Tomography in the Assessment of Patients with 43 Cardiovascular Disease. 2010, 809-824 Duty-Bound: Rational Foundations of Clinical Strategies for Prevention of Cardiovascular Events. 42 2011, 587-596 Cardiovascular Stress Testing. 2012, 79-103 41 PET in Cardiovascular Diseases. 2013, 255-331 40 Evaluation of Cardiovascular Disease in Patients with Diabetes Mellitus Using Myocardial Perfusion 0.1 39 SPECT. Journal of Korean Diabetes, 2012, 13, 191 38 Noninvasive Diagnosis of Coronary Artery Disease in Women. 2014, 61-81 Cardiovascular System. 2014, 157-189 37 Perfusion Measurements of the Myocardium: Radionuclide Methods and Related Techniques. 2014, 1-89 36 What are the niches for SPECT versus PET versus MRI?. Developments in Cardiovascular Medicine, 35 **1998**, 311-322 Heart Scan. 1998, 239-284 34 Nuclear Cardiology 2: Myocardial Perfusion, Metabolism, Infarction, and Receptor Imaging. 2015, 463-528 Adding CT Measurements of Coronary Artery Calcification to Nuclear Myocardial Perfusion Imaging 32 for Risk Stratification. 2015, 225-239 Serial stress myocardial perfusion imaging in patients with percutaneous coronary intervention: a 31 retrospective single-center cohort study. Journal of the Japanese Coronary Association, 2015, 21, 20-27 Myocardial Perfusion by Dual Energy CT. 2015, 195-230 30 The use of pre-test and post-test probability values as criteria before selecting patients to undergo coronary angiography in patients who have ischemic findings on myocardial perfusion scintigraphy. 0.8 29 Anatolian Journal of Cardiology, 2016, 16, 512-519 Stress Dual-Energy Computed Tomography-Myocardial Perfusion Imaging to Identify Coronary Artery Stenoses Causing Ischemia: A Direct Comparison between Invasive Coronary Angiography 28 0.2 0 and Cardiac Magnetic Resonance-Myocardial Perfusion Imaging. Cardiovascular Imaging Asia, 2017, Current Status of Myocardial Perfusion Imaging in Taiwan. Annals of Nuclear Cardiology, 2019, 5, 111-1130.3 27

26	Preoperative nuclear stress testing in the very old patient population. <i>World Journal of Cardiology</i> , 2020 , 12, 210-219	2.1	
25	Cardiovascular event risk estimated by myocardial perfusion SPECT combined with clinical data. <i>Journal of Cardiology</i> , 2021 ,	3	1
24	The Role of Imaging. 2020 , 633-658		
23	Testing Real-World Application of Appropriateness Criteria of Single Photon Emission Computed Tomography (SPECT) In Two Egyptian Hospitals. <i>Egyptian Heart Journal</i> , 2020 , 72, 3	1.3	
22	Nuclear Imaging with Exercise Testing. 2009 , 121-142		
21	Basis of Cardiac Imaging 2: Myocardial Perfusion, Metabolism, Infarction, and Receptor Imaging in Coronary Artery Disease and Congestive Heart Failure. 2006 , 352-394		
20	Value of C-reactive Protein/Albumin Ratio for Predicting Ischemia in Myocardial Perfusion Scintigraphy. <i>Molecular Imaging and Radionuclide Therapy</i> , 2020 , 29, 112-117	0.8	
19	A myocardial extraction method using deep learning for 99mTc myocardial perfusion SPECT images: A basic study to reduce the effects of extra-myocardial activity <i>Computers in Biology and Medicine</i> , 2021 , 141, 105164	7	1
18	Potential Dynamic Imaging of Stationary Dual-Head SPECT with Multi-Pinhole Collimators in Human. 2020 ,		
17	Value of C-reactive Protein/Albumin Ratio for Predicting Ischemia in Myocardial Perfusion Scintigraphy. <i>Molecular Imaging and Radionuclide Therapy</i> , 2020 , 29, 112-117	0.8	1
16	Stable Ischemic Heart Disease and Approach to Chronic Chest Pain. 2022, 74-101		
15	Stress Testing, Nuclear Imaging, Coronary CT Angiography, Cardiac MRI, Cardiopulmonary Exercise Testing. 2022 , 714-732		
14	The application of artificial intelligence in nuclear cardiology Annals of Nuclear Medicine, 2022, 36, 111	2.5	1
13	Proceedings of the 4th Invitational Wintergreen Conference. Wintergreen, Virginia, USA. July 12-14, 1998. Abstracts. <i>Journal of Nuclear Cardiology</i> , 1999 , 6, 93-155	2.1	9
12	Nuclear Cardiology in Women and Underrepresented Minority Populations <i>Current Cardiology Reports</i> , 2022 , 1	4.2	
11	Ischemia Trial: Does the Cardiology Community Need to Pivot or Continue Current Practices?. Current Cardiology Reports,	4.2	
10	Clinical Validation of Japanese Normal Myocardial Perfusion Imaging Databases Using Semi-conductor Gamma Camera (D-SPECT). <i>Annals of Nuclear Cardiology</i> , 2022 ,	0.3	
9	Circulatory System (Cardiovascular and Lymphatic Systems). 2022 , 323-383		

CITATION REPORT

1	Myocardial Ischemic Disease. 2013 , 464-489	O
2	The association of catestatin and endocan with the effects of cardiac shock wave therapy: Biomarker sub-study of the randomized, sham procedure-controlled trial. 10,	O
3	Clinical Trials for the Diagnosis and Management of Stable Ischaemic Heart Disease: Context, Status, and Future Implications. 112-121	o
4	The chest pain guidelines revisited: cherry picking from the frequentist tree.	O
5	Determinaciß del nacie de calcio coronario como predictor de isquemia en pacientes del Centro Màico ABC. 2022 , 67, 172-176	O
6	Economic evaluation of single-photon emission-computed tomography versus stress echocardiography in stable chest pain patients. 2022 , 12,	О
7	Sex Difference Among Patients Undergoing Semiquantitative Tc99m Sesta Mibi Myocardial Scintigraphy as a Prognostic Indicator.	
8	Monte Carlo Simulation and Reconstruction: Assessment of Myocardial Perfusion Imaging of Tracer Dynamics With Cardiac Motion Due to Deformation and Respiration Using Gamma Camera With Continuous Acquisition. Frontiers in Cardiovascular Medicine, 9,	5.4