

# Anthropometric and Metabolic Risk Factors for ESRD A Large Population-Based Cohort Study in Austria

PLoS ONE

11, e0161376

DOI: [10.1371/journal.pone.0161376](https://doi.org/10.1371/journal.pone.0161376)

Citation Report

#	ARTICLE	IF	CITATIONS
1	Long-term risk for end-stage kidney disease and death in a large population-based cohort. Scientific Reports, 2018, 8, 7729.	1.6	5
2	Diabetic kidney disease in the elderly: prevalence and clinical correlates. BMC Geriatrics, 2018, 18, 38.	1.1	47
3	Cigarette smoking as a risk factor for diabetic nephropathy: A systematic review and meta-analysis of prospective cohort studies. PLoS ONE, 2019, 14, e0210213.	1.1	23
4	Gamma-glutamyl transferase variability can predict the development of end-stage of renal disease: a nationwide population-based study. Scientific Reports, 2020, 10, 11668.	1.6	11
5	The Triglyceride-Glucose Index and Obesity-Related Risk of End-Stage Kidney Disease in Austrian Adults. JAMA Network Open, 2021, 4, e212612.	2.8	36
6	Smoking and the risk of diabetic nephropathy in patients with type 1 and type 2 diabetes: a meta-analysis of observational studies. Oncotarget, 2017, 8, 93209-93218.	0.8	20
7	Impact of Obesity in Kidney Diseases. Nutrients, 2021, 13, 4482.	1.7	29
8	A Resampling Method to Improve the Prognostic Model of End-Stage Kidney Disease: A Better Strategy for Imbalanced Data. Frontiers in Medicine, 2022, 9, 730748.	1.2	5
9	Cardiovascular Disease Risk Reduction and Body Mass Index. Current Hypertension Reports, 2022, 24, 535-546.	1.5	6
10	Obesity and renal disease: Benefits of bariatric surgery. Frontiers in Medicine, 0, 10, .	1.2	1