

CITATION REPORT

List of articles citing

Analysis of the prevalence and associated factors of overactive bladder in adult Korean men

DOI: 10.1371/journal.pone.0175641
PLoS ONE, 2017, 12, e0175641.

Source: <https://exaly.com/paper-pdf/87044093/citation-report.pdf>

Version: 2024-04-23

This report has been generated based on the citations recorded by exaly.com for the above article. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

#	Paper	IF	Citations
12	Gait speed and overactive bladder in the healthy community-dwelling super elderly: The Sukagawa Study. <i>Neurourology and Urodynamics</i> , 2019 , 38, 2324-2332	2.3	10
11	Health-related quality of life in adult males with lower urinary tract symptoms. <i>Quality of Life Research</i> , 2019 , 28, 2419-2428	3.7	6
10	Mirabegron as a treatment for overactive bladder symptoms in men (MIRACLE study): Efficacy and safety results from a multicenter, randomized, double-blind, placebo-controlled, parallel comparison phase IV study. <i>Neurourology and Urodynamics</i> , 2019 , 38, 295-304	2.3	17
9	The impact of overactive bladder on health-related quality of life in Korea: based on the results of a Korean Community Health Survey. <i>Quality of Life Research</i> , 2021 , 30, 1017-1024	3.7	0
8	Predictive Factors of De Novo Overactive Bladder After Radical Prostatectomy in Patients With Clinically Localized Prostate Cancer: A Prospective Observational Study. <i>The Korean Journal of Urological Oncology</i> , 2021 , 19, 109-116	0.1	0
7	Überaktive Blase (OAB) beim Mann. <i>Urologie in Der Praxis</i> , 2021 , 23, 97-103	0.1	
6	Analysis of the prevalence of and factors associated with overactive bladder in adult Korean women. <i>PLoS ONE</i> , 2017 , 12, e0185592	3.7	11
5	Seasonal Variation of Drug Prescription Rate for Overactive Bladder in Men Based on National Health Insurance Claims Data, 2012-2016. <i>International Neurourology Journal</i> , 2020 , 24, 278-285	2.6	1
4	Non-linear Relationship Between Body Mass Index and Lower Urinary Tract Symptoms in Korean Males. <i>Journal of Preventive Medicine and Public Health</i> , 2019 , 52, 147-153	3.7	0
3	Monotherapy with mirabegron had a better tolerance than the anticholinergic agents on overactive bladder: A systematic review and meta-analysis. <i>Medicine (United States)</i> , 2021 , 100, e27469	1.8	
2	Monotherapy with mirabegron had a better tolerance than the anticholinergic agents on overactive bladder. <i>Medicine (United States)</i> , 2021 , 100, e27469	1.8	
1	Physical activity and risk factors in subjects with overactive bladder. 2022 , 22, 32-39		0