Effects of a high-intensity pulmonary rehabilitation proventilation/carbon dioxide output slope during exercise undergoing lung resection for non-small cell lung cance

Jornal Brasileiro De Pneumologia 45, e20180132

DOI: 10.1590/1806-3713/e20180132

Citation Report

#	Article	IF	CITATIONS
1	COVID-19 and the elderly: insights into pathogenesis and clinical decision-making. Aging Clinical and Experimental Research, 2020, 32, 1599-1608.	1.4	277
2	Implications of the Adiponectin System in Non-Small Cell Lung Cancer Patients: A Case-Control Study. Biomolecules, 2020, 10, 926.	1.8	15
3	Associations between pretreatment physical performance tests and treatment complications in patients with non-small cell lung cancer: A systematic review. Critical Reviews in Oncology/Hematology, 2021, 158, 103207.	2.0	7
4	Intercept of minute ventilation versus carbon dioxide output relationship as an index of ventilatory in chronic obstructive pulmonary disease. Journal of Thoracic Disease, 2021, 13, 1553-1563.	0.6	1
5	Food, Nutrition, Physical Activity and Microbiota: Which Impact on Lung Cancer?. International Journal of Environmental Research and Public Health, 2021, 18, 2399.	1.2	8
6	Exertional ventilation/carbon dioxide output relationship in COPD: from physiological mechanisms to clinical applications. European Respiratory Review, 2021, 30, 200190.	3.0	6
7	Pulmonary rehabilitation: various diseases, many approaches, and multiple questions. Jornal Brasileiro De Pneumologia, 2019, 45, e20190351.	0.4	0
8	Elderly with COPD: comoborbitidies and systemic consequences. Journal of Gerontology and Geriatrics, 2021, 69, 32-44.	0.2	6
9	Ventilatory efficiency slope is associated with cardiopulmonary complications after thoracoscopic anatomical lung resection. Interactive Cardiovascular and Thoracic Surgery, 2022, 35, .	0.5	2
10	Evolving concepts in COPD and lung cancer: a narrative review. Minerva Medica, 2022, 113, .	0.3	11