

Overweight is associated with allergy in school children Japan

Journal of Medical Investigation

52, 33-40

DOI: [10.2152/jmi.52.33](https://doi.org/10.2152/jmi.52.33)

Citation Report

#	ARTICLE	IF	CITATIONS
2	Levels of Th2 chemokine CCL27 in adolescents: No impact of obesity. <i>Pediatric Allergy and Immunology</i> , 2009, 20, 507-508.	1.1	0
3	U-Shaped Association between Body Mass Index and the Prevalence of Wheeze and Asthma, but not Eczema or Rhinoconjunctivitis: The Ryukyus Child Health Study. <i>Journal of Asthma</i> , 2011, 48, 804-810.	0.9	29
4	Food Allergy: Temporal Trends and Determinants. <i>Current Allergy and Asthma Reports</i> , 2012, 12, 346-372.	2.4	46
5	Influence of health behaviours on the incidence of infection and allergy in adolescents: the AFINOS cross-sectional study. <i>BMC Public Health</i> , 2014, 14, 19.	1.2	11
6	Association of atopic dermatitis with being overweight and obese: A systematic review and metaanalysis. <i>Journal of the American Academy of Dermatology</i> , 2015, 72, 606-616.e4.	0.6	225
7	Contribution of early nutrition on the development of malnutrition and allergic diseases in the first year of life: a study protocol for the Mother and Infant Cohort Study (MICOS). <i>BMC Pediatrics</i> , 2018, 18, 233.	0.7	11
8	Primary Prevention of Food Allergy – Environmental Protection beyond Diet. <i>Nutrients</i> , 2021, 13, 2025.	1.7	10
9	Elevated Body Mass Index is associated with severity of allergic rhinitis: results from a cross sectional study. <i>Australasian Medical Journal</i> , 2008, 1, .	0.1	1
10	Obesity and Lung Health in Children. , 2013, , 217-239.		1
11	Association of body mass index with asthma, allergy rhinitis, and atopic dermatitis among adolescents in Incheon, South Korea. <i>Allergy Asthma & Respiratory Disease</i> , 2014, 2, 243.	0.3	6