

Molecular evolution and phylogenetic implications of intron sequences of Ribosomal DNA in Winteraceae

DOI: [10.2307/2445752](https://doi.org/10.2307/2445752)

Citation Report

#	ARTICLE	IF	CITATIONS
1	Radiation of the endemic genus <i>Dendroseris</i> (Asteraceae) on the Juan Fernandez Islands: evidence from sequences of the ITS regions of nuclear ribosomal DNA. <i>American Journal of Botany</i> , 1994, 81, 1494-1501.	0.8	161
2	DISCORDANCE BETWEEN NUCLEAR AND CHLOROPLAST PHYLOGENIES IN THE <i>HEUCHERA</i> GROUP (SAXIFRAGACEAE). <i>Evolution; International Journal of Organic Evolution</i> , 1995, 49, 727-742.	1.1	318
3	Evoluzione molecolare delle sequenze nucleotidiche degli spaziatori interni dei geni per l'RNA ribosomale in Angiosperme, Gimnosperme e Pteridofite. <i>Giornale Botanico Italiano (Florence, Italy)</i> ; Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 342	0.8	10
4	Phylogeny of <i>Corydalis</i> , ITS and morphology. , 1995, , 183-188.		12
5	The prospects for biological control of the free-living stages of nematode parasites of livestock. <i>International Journal for Parasitology</i> , 1996, 26, 915-925.	1.3	50
6	HOMOPLASY CONNECTIONS AND DISCONNECTIONS: GENES AND SPECIES, MOLECULES AND MORPHOLOGY. , 1996, , 37-66.		43
7	Evolution and biogeography of New Zealand <i>Anaphalis</i> (Asteraceae: Gnaphalieae) inferred from rDNA sequences. <i>New Zealand Journal of Botany</i> , 1997, 35, 441-449.	0.8	24
8	A phylogenetic study of <i>Polygonum sect. Tovar</i> (Polygonaceae) based on ITS sequences of nuclear ribosomal DNA. <i>Journal of Plant Biology</i> , 1997, 40, 47-52.	0.9	13
9	ITS sequences and speciation on far eastern <i>Indigofera</i> (Leguminosae). <i>Journal of Plant Research</i> , 1997, 110, 339-346.	1.2	13
10	Phylogenetic Relationships in the Papilionoideae (Family Leguminosae) Based on Nucleotide Sequences of cpDNA (rbcL) and ncDNA (ITS 1 and 2). <i>Molecular Phylogenetics and Evolution</i> , 1997, 8, 65-88.	1.2	130
11	Bamboozled Again! Inadvertent Isolation of Fungal rDNA Sequences from Bamboos (Poaceae:); Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 342	1.2	80
12	Molecular Evolution of the Internal Transcribed Spacers (ITS1 and ITS2) and Phylogenetic Relationships among Species of the Family Cucurbitaceae. <i>Molecular Phylogenetics and Evolution</i> , 1998, 9, 204-219.	1.2	121
13	The phylogeny and biogeography of <i>Gentiana L. sect. Cimnialis</i> (Adans.) Dumort.: A historical interpretation of distribution ranges in the European high mountains. <i>Perspectives in Plant Ecology, Evolution and Systematics</i> , 1998, 1, 121-135.	1.1	57
14	Molecular evidence for an African origin of the Hawaiian endemic <i>Hesperomannia</i> (Asteraceae). <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1998, 95, 15440-15445.	3.3	82
15	Phylogenetic relationships among <i>Fagopyrum</i> species revealed by the nucleotide sequences of the ITS region of the nuclear rRNA gene.. <i>Genes and Genetic Systems</i> , 1998, 73, 201-210.	0.2	61
16	New <i>Fagopyrum</i> species revealed by morphological and molecular analyses.. <i>Genes and Genetic Systems</i> , 1998, 73, 85-94.	0.2	29
17	The importance of dispersal and recent speciation in the flora of New Zealand. <i>Journal of Biogeography</i> , 1999, 26, 1323-1325.	1.4	51
18	Phylogeography of North American populations of the moss species <i>Hylocomium splendens</i> based on the nucleotide sequence of internal transcribed spacer 2 of nuclear ribosomal DNA. <i>Molecular Ecology</i> , 1999, 8, 1037-1042.	2.0	67

#	ARTICLE	IF	CITATIONS
19	Molecular evidence for the phylogenetic position of <i>Hanabusaya asiatica</i> Nakai (Campanulaceae), an endemic species in Korea. <i>Journal of Plant Biology</i> , 1999, 42, 168-173.	0.9	8
20	A phylogenetic and biogeographic study of <i>Hamamelis</i> (Hamamelidaceae), an eastern Asian and eastern North American disjunct genus. <i>Biochemical Systematics and Ecology</i> , 1999, 27, 55-66.	0.6	49
21	Internal Transcribed Spacer Sequence Phylogeny of <i>Crambe</i> L. (Brassicaceae): Molecular Data Reveal Two Old World Disjunctions. <i>Molecular Phylogenetics and Evolution</i> , 1999, 11, 361-380.	1.2	76
22	ITS Sequence Data Resolve Higher Level Relationships Among the Eucalypts. <i>Molecular Phylogenetics and Evolution</i> , 1999, 12, 215-223.	1.2	68
23	Phylogeny and Evolution of Orchid and Allied Genera Based on ITS DNA Variation: Morphological Gaps and Molecular Continuity. <i>Molecular Phylogenetics and Evolution</i> , 1999, 13, 67-76.	1.2	125
24	Phylogenetic Relationships of the Silver Saxifrages (<i>Saxifraga</i> , Sect. <i>Ligulatae</i> Haworth): Implications for the Evolution of Substrate Specificity, Life Histories, and Biogeography. <i>Molecular Phylogenetics and Evolution</i> , 1999, 13, 536-555.	1.2	80
25	Study of the Evolutionary Relationships among <i>Limonium</i> Species (Plumbaginaceae) Using Nuclear and Cytoplasmic Molecular Markers. <i>Molecular Phylogenetics and Evolution</i> , 2000, 14, 232-249.	1.2	56
26	Sphaeriid and Corbiculid Clams Represent Separate Heterodont Bivalve Radiations into Freshwater Environments. <i>Molecular Phylogenetics and Evolution</i> , 2000, 14, 75-88.	1.2	184
27	Phylogeny of Taxaceae and Cephalotaxaceae Genera Inferred from Chloroplast <i>matK</i> Gene and Nuclear rDNA ITS Region. <i>Molecular Phylogenetics and Evolution</i> , 2000, 14, 353-365.	1.2	84
28	Ribosomal DNA Evidence and Disjunctions of Western American Portulacaceae. <i>Molecular Phylogenetics and Evolution</i> , 2000, 15, 419-439.	1.2	37
29	Xylem sap flow and stem hydraulics of the vesselless angiosperm <i>Drimys granadensis</i> (Winteraceae) in a Costa Rican elfin forest. <i>Plant, Cell and Environment</i> , 2000, 23, 1067-1077.	2.8	54
30	The classification and geography of the flowering plants: Dicotyledons of the class Angiospermae. <i>Botanical Review</i> , The, 2000, 66, 441-647.	1.7	136
31	Testing relative evolutionary rates and estimating divergence times among six genera of Rhizophoraceae using cpDNA and nrDNA sequences. <i>Science Bulletin</i> , 2000, 45, 1011-1015.	1.7	13
32	Genome evolution in polyploids. , 2000, , 225-249.		107
33	Variation along ITS markers across strains of <i>Fibrocapsa japonica</i> (Raphidophyceae) suggests hybridisation events and recent range expansion. <i>Journal of Sea Research</i> , 2001, 46, 213-222.	0.6	31
34	Ribosomal DNA-ITS sequence polymorphism in the sugarcane rust, <i>Puccinia kuehnii</i> . <i>Mycoscience</i> , 2001, 42, 447-453.	0.3	14
35	Phylogeny of the Celastraceae Inferred from 26S Nuclear Ribosomal DNA, Phytochrome B, <i>rbcl</i> , <i>atpB</i> , and Morphology. <i>Molecular Phylogenetics and Evolution</i> , 2001, 19, 353-366.	1.2	89
36	Inferring the History of the Polyploid <i>Silene aegaea</i> (Caryophyllaceae) Using Plastid and Homoeologous Nuclear DNA Sequences. <i>Molecular Phylogenetics and Evolution</i> , 2001, 20, 474-481.	1.2	190

#	ARTICLE	IF	CITATIONS
37	The Tapetum in Basal Angiosperms: Early Diversity. <i>International Journal of Plant Sciences</i> , 2001, 162, 375-392.	0.6	75
38	Evolutionary trends in Winteraceae pollen. <i>Grana</i> , 2002, 41, 4-9.	0.4	14
39	Phylogeographic Congruence Between mtDNA and rDNA ITS Markers in Brown Trout. <i>Molecular Biology and Evolution</i> , 2002, 19, 2161-2175.	3.5	80
40	Plant dispersal N.E.W.S from New Zealand. <i>Trends in Ecology and Evolution</i> , 2002, 17, 514-520.	4.2	184
41	No genetic substructuring in <i>Artemisia</i> subgenus <i>Tridentatae</i> despite strong ecotypic subspecies selection. <i>Biochemical Systematics and Ecology</i> , 2002, 30, 579-593.	0.6	5
42	Three nonorthologous ITS1 types are present in a polypore fungus <i>Trichaptum abietinum</i> . <i>Molecular Phylogenetics and Evolution</i> , 2002, 23, 112-122.	1.2	76
43	Systematics of the tribe Podalyrieae (Fabaceae) based on DNA, morphological and chemical data. <i>Botanical Journal of the Linnean Society</i> , 2002, 139, 159-170.	0.8	18
44	PHYLOGENY OF GAERTNERA LAM. (RUBIACEAE) BASED ON MULTIPLE DNA MARKERS: EVIDENCE OF A RAPID RADIATION IN A WIDESPREAD, MORPHOLOGICALLY DIVERSE GENUS. <i>Evolution; International Journal of Organic Evolution</i> , 2002, 56, 42-57.	1.1	112
45	HARDLY A RELICT: FREEZING AND THE EVOLUTION OF VESSELLESS WOOD IN WINTERACEAE. <i>Evolution; International Journal of Organic Evolution</i> , 2002, 56, 464-478.	1.1	76
46	The Genus <i>Clusia</i> L.: Molecular Evidence for Independent Evolution of Photosynthetic Flexibility. <i>Plant Biology</i> , 2002, 4, 86-93.	1.8	27
47	The effects of increasing genetic distance on alignment of, and tree construction from, rDNA internal transcribed spacer sequences. <i>Molecular Phylogenetics and Evolution</i> , 2003, 26, 444-451.	1.2	42
48	ITS secondary structure derived from comparative analysis: implications for sequence alignment and phylogeny of the Asteraceae. <i>Molecular Phylogenetics and Evolution</i> , 2003, 29, 216-234.	1.2	141
49	Determination of the genetic relationship of <i>Dendrobium</i> species (Orchidaceae) in Taiwan based on the sequence of the internal transcribed spacer of ribosomal DNA. <i>Scientia Horticulturae</i> , 2004, 101, 315-325.	1.7	32
50	Nuclear ribosomal DNA internal transcribed spacer 1 (ITS1) in <i>Picea</i> (Pinaceae): sequence divergence and structure. <i>Molecular Phylogenetics and Evolution</i> , 2005, 35, 165-185.	1.2	53
51	The internal transcribed spacer of nuclear ribosomal DNA in the gymnosperm <i>Gnetum</i> . <i>Molecular Phylogenetics and Evolution</i> , 2005, 36, 581-597.	1.2	78
52	Nuclear ribosomal DNA sequence variation and evolution of spotted marsh-orchids (<i>Dactylorhiza</i>) Tj ETQq1 1 0.784314 rgBT JOverloc 1.2 33	1.2	33
53	CLIMATE AND LIFE-HISTORY EVOLUTION IN EVENING PRIMROSES (OENOTHERA, ONAGRACEAE): A PHYLOGENETIC COMPARATIVE ANALYSIS. <i>Evolution; International Journal of Organic Evolution</i> , 2005, 59, 1914-1927.	1.1	75
54	Molecular identification of Korean ginseng by amplification refractory mutation system-PCR. <i>Food Research International</i> , 2006, 39, 568-574.	2.9	42

#	ARTICLE	IF	CITATIONS
55	A Nuclear Ribosomal DNA Phylogeny of <i>Acer</i> Inferred with Maximum Likelihood, Splits Graphs, and Motif Analysis of 606 Sequences. <i>Evolutionary Bioinformatics</i> , 2006, 2, 117693430600200.	0.6	51
56	Phylogenetic relationships of <i>Rhododendron</i> section <i>Vireya</i> (Ericaceae) inferred from the ITS nrDNA region. <i>Australian Systematic Botany</i> , 2006, 19, 329.	0.3	9
57	Sequence variations of the first ribosomal internal transcribed spacer of <i>Penaeus</i> species in Thailand. <i>Journal of Experimental Marine Biology and Ecology</i> , 2006, 331, 64-73.	0.7	11
58	A survey of nuclear ribosomal internal transcribed spacer substitution rates across angiosperms: an approximate molecular clock with life history effects. <i>BMC Evolutionary Biology</i> , 2006, 6, 36.	3.2	291
59	The evolutionary history of <i>Sium</i> sensu lato (Apiaceae): dispersal, vicariance, and domestication as inferred from ITS rDNA phylogeny. <i>American Journal of Botany</i> , 2006, 93, 747-761.	0.8	45
60	Evolution of <i>Piperales</i> matK gene and trnK intron sequence data reveal lineage specific resolution contrast. <i>Molecular Phylogenetics and Evolution</i> , 2007, 42, 477-497.	1.2	127
61	Nuclear ribosomal pseudogenes resolve a corroborated monophyly of the eucalypt genus <i>Corymbia</i> despite misleading hypotheses at functional ITS paralogs. <i>Molecular Phylogenetics and Evolution</i> , 2007, 44, 752-764.	1.2	47
62	Relationships of <i>Anagallis foemina</i> and <i>A. arvensis</i> (Myrsinaceae): New insights inferred from DNA sequence data. <i>Molecular Phylogenetics and Evolution</i> , 2007, 45, 971-980.	1.2	10
63	Variation and similarities in pollen features in some basal angiosperms, with some taxonomic implications. <i>Plant Systematics and Evolution</i> , 2007, 263, 59-75.	0.3	14
64	Phylogeny of <i>Bonatea</i> (Orchidaceae: Habenariinae) based on molecular and morphological data. <i>Plant Systematics and Evolution</i> , 2007, 263, 253-268.	0.3	13
65	A phylogenetic analysis of Dipsacaceae based on four DNA regions. <i>Plant Systematics and Evolution</i> , 2009, 279, 69-86.	0.3	26
66	Class Magnoliopsida (Dicotyledons)., 2009, , 7-588.		3
67	Chemical constituents and DNA sequence analysis of a psychotropic herbal product. <i>Forensic Toxicology</i> , 2010, 28, 77-83.	1.4	19
68	Evolution And Diversification Of Brook Weeds (<i>Samolus</i> , Samolaceae, Ericales). <i>International Journal of Plant Sciences</i> , 2011, 172, 250-266.	0.6	5
69	Phylogenetic Analysis of <i>Phyllanthus</i> Species. <i>Traditional Herbal Medicines for Modern Times</i> , 2011, , 71-95.	0.1	0
70	Internal transcribed spacer (ITS) and genetic variations in <i>Lilium</i> native to Korea. <i>Horticulture Environment and Biotechnology</i> , 2011, 52, 502-510.	0.7	13
71	Genetic variability and relationships among interspecific hybrid cultivars and parental species of <i>Paphiopedilum</i> via ribosomal DNA sequence analysis. <i>Plant Systematics and Evolution</i> , 2012, 298, 1897-1907.	0.3	5
72	Genetic and chemical components analysis of <i>Papaver setigerum</i> naturalized in Korea. <i>Forensic Science International</i> , 2012, 222, 387-393.	1.3	13

#	ARTICLE	IF	CITATIONS
73	Assessment of phylogenetic inter-relationships in the genus <i>Cymbidium</i> (Orchidaceae) based on internal transcribed spacer region of rDNA. <i>Gene</i> , 2012, 495, 10-15.	1.0	25
74	Using nuclear gene data for plant phylogenetics: Progress and prospects. <i>Molecular Phylogenetics and Evolution</i> , 2012, 65, 774-785.	1.2	101
75	DNA barcodes reveal high levels of morphological plasticity among <i>Rhododendron</i> species (Ericaceae) in Taiwan. <i>Biochemical Systematics and Ecology</i> , 2012, 40, 169-177.	0.6	7
76	Evolution of the <i>Portulaca oleracea</i> L. aggregate in Egypt on molecular and phenotypic levels revealed by morphology, inter-simple sequence repeat (ISSR) and 18S rDNA gene sequence markers. <i>Flora: Morphology, Distribution, Functional Ecology of Plants</i> , 2013, 208, 464-477.	0.6	19
77	Reprint of: Using nuclear gene data for plant phylogenetics: Progress and prospects. <i>Molecular Phylogenetics and Evolution</i> , 2013, 66, 539-550.	1.2	52
78	Reticulate evolution in <i>Ranunculus cantonensis</i> polyploid complex and its allied species. <i>Plant Systematics and Evolution</i> , 2013, 299, 603-610.	0.3	3
79	Interglacial genetic diversification of <i>Moussonia deppeana</i> (<sc>G</sc>esneriaceae), a hummingbirdâ€pollinated, cloud forest shrub in northern <sc>M</sc>esoamerica. <i>Molecular Ecology</i> , 2014, 23, 4119-4136.	2.0	25
80	Convergence of anti-bee pollination mechanisms in the Neotropical plant genus <i>Drymonia</i> (Gesneriaceae). <i>Evolutionary Ecology</i> , 2015, 29, 355-377.	0.5	32
81	Comparative Characterization of Ribosomal DNA Regions in Different <i>Agave</i> Accessions with Economical Importance. <i>Plant Molecular Biology Reporter</i> , 2015, 33, 2014-2029.	1.0	6
82	How just a few makes a lot: Speciation via reticulation and apomixis on example of European brambles (<i>Rubus</i> subgen. <i>Rubus</i> , Rosaceae). <i>Molecular Phylogenetics and Evolution</i> , 2015, 89, 13-27.	1.2	81
83	Sequence analysis of the internal transcribed spacer (ITS) region of the nuclear ribosomal DNA (nrDNA) <i>Chrysanthemum</i> species in Korea. <i>Horticulture Environment and Biotechnology</i> , 2015, 56, 44-53.	0.7	9
84	Intercontinental long-distance dispersal of Canellaceae from the New to the Old World revealed by a nuclear single copy gene and chloroplast loci. <i>Molecular Phylogenetics and Evolution</i> , 2015, 84, 205-219.	1.2	29
85	A mistletoe tale: postglacial invasion of <i>Psittacanthus schiedeana</i> (Loranthaceae) to Mesoamerican cloud forests revealed by molecular data and species distribution modeling. <i>BMC Evolutionary Biology</i> , 2016, 16, 78.	3.2	63
86	In and out of the Qinghaiâ€Tibet Plateau: divergence time estimation and historical biogeography of the large arcticâ€alpine genus <i>Saxifraga</i> L.. <i>Journal of Biogeography</i> , 2017, 44, 900-910.	1.4	117
87	Non-congruent geographic patterns of genetic divergence across European species of <i>Branchinecta</i> (Anostraca: Branchinectidae). <i>Hydrobiologia</i> , 2017, 801, 47-57.	1.0	13
88	On the Evolutionary and Biogeographic History of <i>Saxifraga</i> sect. <i>Trachyphyllum</i> (Gaud.) Koch (<i>Saxifragaceae</i> Juss.). <i>PLoS ONE</i> , 2013, 8, e69814.	1.1	25
89	Leaf morphological and genetic divergence in populations of <i>Drimys</i> (Winteraceae) in Chile. <i>Genetics and Molecular Research</i> , 2012, 11, 229-243.	0.3	7
90	Molecular Identification of Korean Mountain Ginseng Using an Amplification Refractory Mutation System (ARMS). <i>Journal of Ginseng Research</i> , 2010, 34, 41-46.	3.0	16

#	ARTICLE	IF	CITATIONS
91	Reproductive ecology and isolation of <i>Psittacanthus calyculatus</i> and <i>P. auriculatus</i> mistletoes (Loranthaceae). <i>PeerJ</i> , 2016, 4, e2491.	0.9	22
92	Phylogenetic Study of Korean <i>Chrysosplenium</i> Based on nrDNA ITS Sequences. <i>Korean Journal of Plant Resources</i> , 2011, 24, 358-369.	0.2	7
93	Molecular Phylogenetic Studies of Korean <i>Hydrocotyle</i> L.. <i>Korean Journal of Plant Resources</i> , 2012, 25, 490-497.	0.2	5
94	A Phylogenetic Study of Korean <i>Carpesium</i> L. Based on nrDNA ITS Sequences. <i>Korean Journal of Plant Resources</i> , 2012, 25, 96-104.	0.2	3
95	Comparative analysis of nucleotide sequences of 5.8S nuclear ribosomal DNA of 30 plant species. <i>Biopolymers and Cell</i> , 1997, 13, 150-156.	0.1	0
96	An attempt to combine morphological characters and nuclear ribosomal DNA (internal transcribed) Tj ETQq1 1 0.784314 rgBT /Overl 383-389.	0.1	3