

Length-weight, length-length relationships and condition indices of *Carassius auratus* (Actinopterygii: Cyprinidae) from the southern Caspian Sea

Journal of Animal Diversity

2, 56-61

DOI: [10.29252/jad.2020.2.2.6](https://doi.org/10.29252/jad.2020.2.2.6)

Citation Report

#	ARTICLE	IF	CITATIONS
1	Predicting the Trace Element Levels in Caspian Kutum ( <i>Rutilus kutum</i> ) from South of the Caspian Sea Based on Locality, Season and Fish Tissue. <i>Biological Trace Element Research</i> , 2022, 200, 354-363.	1.9	2
2	Evaluation of Histopathological Effect of Roach ( <i>Rutilus rutilus caspicus</i> ) in Exposure to Sub-Lethal Concentrations of Abamectin. <i>Water, Air, and Soil Pollution</i> , 2021, 232, 1.	1.1	25
3	Histological effects of sublethal concentrations of insecticide Lindane on intestinal tissue of grass carp ( <i>Ctenopharyngodon idella</i> ). <i>Veterinary Research Communications</i> , 2021, 45, 373-380.	0.6	22
4	Relationship Between Trace Element Content in the Brain of Bony Fish Species and Their Food Items in the Southwest of the Caspian Sea Due to Anthropogenic Activities. <i>Avicenna Journal of Environmental Health Engineering</i> , 2020, 7, 78-85.	0.3	2
5	A Review on the Effects of Heavy Metals on Aquatic Animals. <i>Journal of Biomedical Research &amp; Environmental Sciences</i> , 2021, 2, 865-869.	0.1	13
6	Heavy Metals in the Fish <i>Tenulosa ilisha</i> Hamilton, 1822 in the Padma-Meghna River Confluence: Potential Risks to Public Health. <i>Toxics</i> , 2021, 9, 341.	1.6	9
7	Estimating length-weight, length-length relationships, and condition factor of eight fish species, a case study of Bashar River, Tigris drainage (Iran). <i>Su Üzerindeki Dergisi</i> , 2022, 39, 332-337.	0.1	0